

University of Groningen

Genetics of manta and devil rays

Poortvliet, Marloes

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Poortvliet, M. (2015). *Genetics of manta and devil rays: Evolution, population genetics and conservation of a group of vulnerable pelagic filter-feeders*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

8

Discussion, Summary and Samenvatting



DISCUSSION

What is in a name?

Over the past 20 years our understanding of evolutionary relationships across the tree of life have drastically changed, with the availability of ever-improving molecular phylogenies and more comprehensive sampling (de Queiroz & Gauthier 1990). At the same time, however, branching relationships among clades have shown themselves to be unstable in many cases. For example, higher-level taxonomic groups are often shown to be paraphyletic (or even polyphyletic), i.e. the recognized taxonomic rank only includes a portion of the descendants. Paraphyly most commonly occurs in the tree where taxon sampling has been poor, for example where an “exemplar” approach has been applied in which one individual is used to represent a species, or one species to represent a genus, or one genus to represent a family, etc. Additionally, cryptic species (those that are morphologically indistinguishable but very different at the molecular level) and conversely, morphologically distinctive taxa that are indistinguishable at the molecular level (based on the sequences employed), can also cause paraphyly.

Paraphyly creates nomenclatural problems in the application of Linnaean classification, because the tree of life is based on identity by descent and the recognition of monophyletic groups. The most commonly held view is that a genus or higher-level group should be monophyletic, i.e., a group should contain all species that share the most recent common ancestor (Henning 1966). Therefore, when paraphyly is discovered and confirmed from various data sources, it should be resolved by making nomenclatural changes to the original Linnaean classification. This can be relatively easy to do and if so, should be carried out. If, however, the changes required by nomenclatural priority, as outlined in the Code of Zoological Nomenclature (CZN), result in numerous changes and greater ambiguity in the taxonomy and recognition of the taxon, then measures can be taken to conserve the Latin binomial as a matter of practicality and legality. Indeed, taxonomic stability is a concept that is positively valued in the CZN, and can, in some instances, take precedence over other principles. The importance of a name is more than esoteric. It provides a lingua franca and a legal status that is especially relevant in conservation and species protections.

Results from Chapter 5 indicate that the genus *Mobula* is paraphyletic with respect to *Manta*. The two *Manta* species form a clade together with *Mobula tarapacana* and *Mobula japanica*, a result that is in agreement with previous studies based on molecular data (Aschliman *et al.* 2012a; Aschliman *et al.* 2012b; Naylor *et al.* 2012) and morphology (Adnet *et al.* 2012; Aschliman *et al.* 2012a; Gonzalez-Isais & Dominguez 2004; Herman *et al.* 2000). Based on the principle of monophyly, *Manta* should therefore be renamed to *Mobula*, since *Mobula* (Rafinesque-Schmaltz 1810) has nomenclatural priority over *Manta* (Bancroft 1831). However, it is necessary to consider the practical and legal aspects of a possible name change. Of paramount importance is, in this respect, the issue of conservation. Conservation is based on formal species names, and changing a species' Latin binomial name can potentially influence its conservation status and, as a result, the conservation effort for these vulnerable species.

Manta rays are currently protected in several countries, including Mexico, Ecuador, the Republic of Maldives and the Philippines. Additionally, *Mobula birostris* is listed on Appendix I and II of the Convention of Migratory Species (CMS), and is protected by EU regulations (no. 43/2014), and *M. birostris*, *Mobula alfredi* and the putative third *Manta* species have received protection under the Convention on International Trade in Endangered Species (CITES) Appendix II. It is difficult to predict the consequences of a name change from *Manta* to *Mobula* on each individual conservation measure applicable to *Manta*. However, a review of taxonomic changes and their impact on conservation effort shows that name changes negatively impact conserva-

tion efforts only when they concern threatened species that have been lumped together with a non-threatened species, thereby effectively removing them from any protective category (Morrison *et al.* 2009). For example, the Idaho springsnail (*Pyrgulopsis idahoensis*) was removed from Endangered Species Act (ESA) after it was shown to be conspecific to the more common Jackson Lake springsnail (Hershler & Liu 2004), resulting in decreased conservation efforts. However, it has also been observed that name changes involving iconic species have little effect on conservation effort, even when they concern the lumping of a threatened species with a non-threatened one (Morrison *et al.* 2009). A case-by-case examination of taxonomic changes and their effect on conservation effort does not necessarily prove that renaming *Manta* to *Mobula* would have no impact on conservation effort. However, the iconic status of manta rays, as well as the fact that both manta rays remain “good” species even if their Latin names are changed, suggests that manta rays will not be negatively impacted if the genus *Manta* ceases to exist.

Species as units of conservation

Knowledge of the full biodiversity of a genus or Family facilitates the identification of appropriate units for conservation and the associated research necessary (Simpfendorfer *et al.* 2011). However, it can also create dilemmas. Based on results from Chapter 5, all of the currently described small (max. DW <130 cm) and morphologically similar mobulid species represent distinct evolutionary lineages: First, *Mobula hypostoma* and *Mobula rochebrunei*, which are distributed in the Western and Eastern Atlantic respectively, diverged around 1.1 Mya, and are most closely related to East Pacific *Mobula munkiana*; Second, *Mobula eregoodootenkee* and *Mobula kuhlii*, two small and morphologically similar mobulid species with overlapping distributions in the Indo-West Pacific, diverged more recently at around 500 Kya. These small mobulids are among the least studied species in the Family. With the exception of *M. munkiana*, almost nothing is known about their biology beyond a taxonomic record. Nevertheless, results from Chapter 5 suggest that they are frequently caught in inshore artisanal fisheries and as bycatch in tuna purse seine fisheries (Croll *et al.* submitted), especially across the West Pacific and Indian Oceans (i.e. *M. eregoodootenkee* and *M. kuhlii*). Population trends are unknown, but based on their (likely) limited reproductive potential and high catch rates throughout their ranges, *M. eregoodootenkee* and *M. munkiana* are listed as ‘Near Threatened’ and *M. rochebrunei* as ‘Vulnerable’. *M. kuhlii* and *M. hypostoma* are listed as ‘Data Deficient’. More comprehensive threat assessments should be conducted for each of these species separately. Where species overlap in the West Pacific and Indian Oceans, genetic markers developed in chapter 3 and 5 could be used to genotype fisheries catches, as well as dried gill rakers, in order to determine catch rates for each individual species.

M. japonica and *Mobula mobular* represent a different case in which nomenclatural changes could actually worsen protection. The small amount of mitochondrial and nuclear sequence divergence (Chapters 5 and 6) between the two species, as well as placement of *M. mobular* within the Indian/Pacific mitogenome clade of *M. japonica* (Figure 2), strongly suggests that these two taxa are conspecific, i.e., constitute only one species. If genetic and morphological analyses of additional *M. mobular* samples (which are essential in this case where only an “exemplar approach” has been applied) confirm this result, *M. japonica* (Müller & Henle 1841); should be renamed to *M. mobular* (Bonnaterre 1788), since the latter has nomenclatural priority over the former. *M. mobular* only occurs in the Mediterranean Sea and surrounding Atlantic Ocean. It is classified as ‘Endangered’ on the IUCN Red List, due to unsustainable levels of bycatch in various fisheries (Notarbartolo di Scara *et al.* 2007) and is currently protected in Croatia and Malta, and through the 1995 Barcelona Convention. Unfortunately, merging *M. japonica*

with *M. mobular* will result in a larger group with a greater range, which in turn might result in decreased conservation efforts for *M. mobular* (Morrison *et al.* 2009). If *M. japanica* is indeed synonymized with *M. mobular*, it is of paramount importance that local and regional protective measures for *M. mobular* stay in place and are further expanded.

Populations as units of conservation

At the species level, knowledge of population structure and the overall pattern of genetic diversity allows for the determination of the appropriate geographic scale for conservation management actions. Two types of conservation units (CUs) are most commonly discussed: evolutionarily significant units (ESUs) and management units (MUs) (Moritz 1994b). Although there are many different definitions of ESUs, the term is generally defined as ‘a population or group of populations that warrant separate management or priority for conservation because of high genetic and ecological distinctiveness’. At a smaller scale, MUs are populations that are demographically independent within one ESU, and which should be managed separately to ensure the long-term persistence of species.

Based on criteria suggested by Moritz (1994) for the resolution of ESUs and MUs based on genetic markers, *M. japanica* constitutes one ESU, i.e. there are no detectable adaptive differences between lineages within the species (Chapter 6). However, the species does contain several demographically independent populations, which constitute separate MUs (i.e. Atlantic Ocean and Indian/Pacific Ocean populations), and hence should be managed separately to ensure the long-term persistence of species. Limited genetic structure also exists between the East Pacific and Indian/West Pacific Oceans, but it is unclear whether the difference is biologically relevant, due to uncertainty about the reproductive behavior of *M. japanica*, as well as a lack of broader geographic representation of samples from the West Pacific.

The role of upwelling in the evolutionary and demographic history of mobulids

In conjunction with genetic data, palaeoceanographic data provides information about the evolutionary history of a species. A review of the relevant literature suggests that upwelling of deep, nutrient rich water and related high primary productivity played an important role in the rise and subsequent diversification of the mobulid lineage, as well as their historical demography. This is plausible, given the mode of foraging of this species group. Although upwelling is not the only source of high productivity in the ocean, it does concentrate resources in a relatively small geographic area, often providing densities of food particles high enough to meet the large energy demands of filter-feeding marine mega fauna (Croll *et al.* 2005). The importance of upwelling is also evident from the current distribution and migration patterns of mobulids, which for a large part overlap with or follow seasonal upwelling (Anderson *et al.* 2011; Croll *et al.* 2012; Graham *et al.* 2012).

Upwelling is not, however, a stable phenomenon through time, but has waxed and waned following changes in the world’s climate throughout the ages. Periods of global cooling have generally been associated with high upwelling intensities (e.g. during the Early and Middle Oligocene), while upwelling has markedly decreased during periods of global warming. Hence, climatic fluctuations have profoundly affected conditions experienced by pelagic marine filter-feeders. In Chapters 5 & 6 I suggest that upwelling has been an important driver of inter- and intra-specific phylogenetic patterns in mobulids:

First, the rise of the mobulid lineage (Chapter 5) around 30 Mya coincided with a period of global cooling (Figure 1), which, together with tectonics-mediated changes in ocean currents, facilitated massive upwelling of nutrient-rich deep waters. This in turn greatly boosted produc-

tivity in tropical and subtropical regions. It is possible that the development of the filter-feeding strategy in mobulids – a key innovation within the Order Batoidae—was mediated by this increase in upwelling, i.e. through ecological opportunity. The upwelling-hypothesis has also been proposed for other large filter-feeders, including mysticeti whales (Fordyce 1980).

Second, later divergences within the mobulid lineage (Chapter 5) largely coincided with periods of global warming (Figure 1), during which upwelling and related primary productivity was generally reduced. This reduction in upwelling would have resulted in the fragmentation of the habitat of filter-feeders, leading to isolation of subpopulations in different regions with limited connectivity between them. If the time of reduced upwelling was sufficiently prolonged, reproductively isolated populations may have diverged enough to evolve into new species. Other (temporary) barriers to dispersal, such as the presence of land-barriers during periods of low sea levels may have further reinforced divergence between species through ecological specialization. Pastene *et al.* (2007) proposed a similar theory for speciation in the minke whales, a temperate baleen filter-feeder.

Third, three of the four mitogenome clades present in *M. japanica* show evidence of demographic expansion during times of increased upwelling during the Pleistocene (Chapter 6). This pattern of expansion is consistent with an overall increase in carrying capacity of existing upwelling regions through increased availability of nutrients, as well as the rise of new upwelling regions. Demographic expansions during the Pleistocene have been observed in various pelagic species and are inferred to be the result of increased upwelling-related primary productivity (Alvarado Bremer *et al.* 2005; Amaral *et al.* 2012; Diaz-Jaimes *et al.* 2010; Duncan *et al.* 2006; Ely *et al.* 2005; Hoelzel *et al.* 2006; Martinez *et al.* 2006; Theisen *et al.* 2008).

How species have responded to past climate fluctuations may provide critical insights about how that species may respond to current climate change (O'Brien *et al.* 2012). Many unknowns remain in projecting the potential effect of climate change on upwelling systems, but based on the IPCC Special Report on Emission Scenario A2, equatorial upwelling systems are projected to decrease by almost 30 % by 2100 (Polovina *et al.* 2011). If correct, such a reduction will have a tremendous impact on mobulids and other filter-feeders that are strongly dependent on the high productivity associated with upwelling. Demographic declines, due to decreased carrying capacity of upwelling regions, as well as major population range shifts to areas with sufficient food availability are projected outcomes.

In conclusion, the high and increasing catch rates in various (bycatch) fisheries, as well as projected contractions of important feeding regions due to climate change pose a formidable threat to the continued existence of this vulnerable group of pelagic elasmobranchs. It is of paramount importance that we act with urgency, before they are lost forever.

Summary

SUMMARY

Mobulids are large pelagic filter-feeders and amongst the most vulnerable of all chondrichthyan species due to slow life histories and extremely low fecundity. Unfortunately, large and expanding fisheries exist in various Asian countries driven by the use of mobulid filter plates in Chinese medicine. Mobulids are also frequently caught as bycatch in tuna purse seine fisheries across their entire range. Due to the low intrinsic growth rate of mobulid populations, even low catch rates can result in significant population declines.

The evolutionary history of mobulid rays has remained enigmatic despite considerable recent interest in mobulid ecology and conservation. This thesis was designed to provide baseline information on mobulid evolution and population genetics, in order to inform mobulid management and better target conservation measures. I used Next Generation Sequencing of entire mitochondrial genomes, conventional Sanger sequencing and supplementary GenBank sequences to reconstruct—for the first time—evolutionary relationships among all currently described mobulid species. I also analyzed intra-specific patterns of genetic variation in one of the most frequently caught species in various fisheries, the circumtropically distributed spinetail devilray, *Mobula japanica*.

The general questions of this thesis were:

1. What are the evolutionary relationships within the Family Mobulidae, and what is the geological timing and pattern of their radiation? What were the probable mechanisms and drivers of mobulid speciation?
2. What is the spatial scale and connectivity of genetically defined populations of the spinetail devil ray, *Mobula japanica*? What were the influences of past climate change on its historical population structure and demography?
3. Is there evidence for sex-biased dispersal in *M. japanica*?
4. What are the main anthropogenic threats to mobulids? How can this group of species be better protected?

The advent of next-generation sequencing (NGS) now makes it possible to cheaply and efficiently obtain large data sets, in comparison with traditional Sanger sequencing or fragment analyses. The application of NGS seems almost endless, and allows for increased resolution in many fields related to biological sciences, including intra- and interspecific studies. For this thesis, the use of mitogenomes resulted in the recovery of highly supported phylogenetic patterns as well as the recovery of four mitogenome clades in *M. japanica*, compared to analyses of individual mitochondrial genes and microsatellites.

Phylogeny of manta and devil rays

Despite the formal separation of mobulids into two genera (i.e. *Manta* and *Mobula*), my research showed that the genus *Mobula* is paraphyletic with the inclusion of *Manta* (Chapter 5). Although paraphyly has been previously suggested based on morphology (Adnet *et al.* 2012; Aschliman *et al.* 2012b; Gonzalez-Isais & Dominguez 2004; Herman *et al.* 2000), parasite evolution (Benz & Deets 1988) and sequence comparisons among a small number of mobulid species (Aschliman *et al.* 2012a; Naylor *et al.* 2012), this is the first time paraphyly has been

confirmed using DNA sequences of all mobulids. My research further showed that mobulids are divided into three distinct clades (Figure 1). Based on a fossil-calibrated phylogenetic analysis, I estimated the rate of mitochondrial evolution in mobulids at $9.93 \times 10^{-9} \text{ S s}^{-1} \text{ y}^{-1}$, which is around an order of magnitude slower than rates found in mammals (Martin *et al.* 1992) and concordant with estimates for other chondrichthyan species (Dudgeon *et al.* 2012). Based on this rate, the Family Mobulidae diverged from its closest relatives the cownose rays (Family *Rhinoptera*) during the Oligocene, around 30 Mya. This divergence marked the transition from a durophagous (shell crushing) to a filter-feeding strategy. Subsequent divergences of species occurred during the Early and Middle Miocene (19-17 Mya), followed by long internal branches and a second interval of divergence during the Pliocene and Pleistocene (3.6 Mya – recent) (Figure 1). I speculate that the waxing and waning of upwelling of deep, nutrient rich water, which directly influences the availability of food, has played a large role in the rise and diversification of the mobulid lineage.

Phylogeography and population genetic structure in *Mobula japonica*

In Chapter 6 I resolved the geographic scale of genetically isolated populations, as well as the demographic history of one of the 11 mobulid species, the circumtropical *M. japonica*. Population genetic structure was assessed using 263 samples from a total of 8 locations from across the Pacific, Indian and Atlantic Oceans, in combination with Sanger sequencing of mitochondrial genes COX1 and NADH5 and 11 microsatellite loci. Results suggested high connectivity across vast oceanic regions. The Atlantic Ocean is genetically differentiated from the Indian and Pacific Oceans (pairwise $F_{ST}(\text{msats})=0.033-0.046$; pairwise $F_{ST}(\text{mtDNA})=0.140-0.167$); however, within the Indian and Pacific Oceans, pairwise comparisons showed an absence of population structure. Only hierarchical AMOVA analyses showed weak genetic structure ($F_{ST}(\text{msats})=0.007$, $F_{ST}(\text{mtDNA})=0.048$), between two East Pacific regions, and the Atlantic and Indian/West Pacific Oceans. There was no evidence for Isolation by Distance.




Figure 1: Divergence time estimates based on analysis of the mitogenome BEAST data set. Point estimates of ages are given above each node and transparent blue horizontal bars denote age (95 % highest posterior density). Branch colors show relative nucleotide substitution rates (blue is slow, purple intermediate, red fast). Red bars below the epoch bar denote periods of global warming and associated low upwelling intensities. Black lines below that denote timing of occurrence of extinct and extant mobulid species based on the fossil record.

A review of population genetic structure found in other chondrichthyans suggested that the dispersal capacity of *M. japanica* is only rivaled by the largest of the pelagic sharks, e.g. whale sharks (Castro *et al.* 2007; Vignaud *et al.* 2014) and basking shark (Hoelzel *et al.* 2006). The similarity in population genetic patterns between these species and *M. japanica* was surprising given that both whale and basking sharks are large pelagic species exhibiting trans-oceanic migration (Eckert *et al.* 2002; Gore *et al.* 2008). *M. japanica* on the other hand displays much more limited seasonal migrations: tagging data suggests migrations between the Gulf of California and adjoining East Pacific Ocean, i.e. across distances of <1500 km. Discrepancies between directly observed dispersal distances and indirectly estimated distances based on genetic data are common. Tagging studies follow individuals for months to several years, whereas population genetic studies integrate time across several generations. It is possible that during El Niño years, when upwelling is generally restricted and productivity in regions frequented by *M. japanica* severely reduced (Escibano *et al.* 2004), the species roams across larger distances in search for high productivity feeding regions, thereby promoting connectivity across much larger regions.

In order to gain more potential resolving power for the detection of cryptic population structure, NGS sequencing of whole mitochondrial genomes was applied. This dramatically increased resolution (11,427 bp) compared to the two-gene mtDNA dataset (COX1/NADH5, 1107 bp). Although population genetic patterns based on 60 mitogenome sequences from four locations were similar to patterns based on the more limited datasets, *M. japanica* individuals were separated into four distinct and well-supported mitogenome clades, with clade membership only loosely determined by geographic location (Figure 2) (Chapter 5). Based on sequence divergence and a mitochondrial clock of $9.93 \times 10^{-9} \text{ S s}^{-1} \text{ y}^{-1}$, the four clades diverged during the Late Pleistocene (126 – 11.7 Kya). I hypothesized that this clade structure is the result of vicariance across physical Pleistocene barriers to dispersal, most notably the cold waters around the Cape of Good Hope during periods of low Agulhas leakage (Peeters 2002), exposure of the Sunda/Sahul shelf during LGM low sea level stands (Clark *et al.* 2009). Both of these barriers have been implicated in the formation of mitochondrial clades in other highly vagile tropical pelagic species, for example tuna, billfishes, cetaceans, and some of the larger pelagic shark species (Amaral *et al.* 2012; Bremer *et al.* 1998; Daly-Engel *et al.* 2012; Duncan *et al.* 2006; Martinez *et al.* 2006; Morin *et al.* 2010; Rosel & Block 1996; Vinas *et al.* 2004). The fact that clade membership was only loosely related to geographic sampling location implies the occurrence of periods of increased connectivity between isolated regions, during which mixing of clades occurred. The four mitogenome clades were indistinguishable based on nuclear microsatellite loci, indicating that individuals are interbreeding and therefore belong to the same species. Finally, three of the four clades showed evidence of demographic expansion, which occurred around 40 Kya and between 10 – 25 Kya. Periodic high upwelling and enhanced primary productivity would have increased the carrying capacity for a pelagic filter-feeder.

Sex-biased dispersal

Many chondrichthyan species display philopatric behavior, i.e. a tendency of individuals to return to their natal site or other adopted locality for breeding (Mayr 1963). In chondrichthyans, philopatry is often sex-biased, with females repeatedly returning to specific sites for breeding or pupping, and males roaming more widely (e.g. blacktip sharks, lemon sharks and white sharks, reviewed by Hueter *et al.* 2004). Such behavior, if present, has important implications for the conservation of the species, as nursery areas constitute essential habitats of a population, and are therefore comparable to natal streams for salmon, or nesting beaches for turtles. No reliable information exists about the occurrence of philopatry in *M. japanica*, although catch




Figure 2: Bayesian phylogenetic tree based on the mitogenome sequences of 60 *Mobula japonica* individuals from Pacific Mexico (PAMX), Pacific Costa Rica (PACR), the Indian Ocean (IO), the East Atlantic Ocean (EAO), one individual of *M. mobular* (MMO), and one individual each of two outgroup species (*M. tarapacana* and *M. birostris*). Branch colors correspond to sampling location. Posterior probabilities (PP) for nodes separating principle clades are shown; all other PP are higher than 0.80 (not shown). Yellow star indicates placement of the outgroup species.

and tagging data suggest that seasonal migration does take place between fixed regions of high primary productivity in the Gulf of California and the surrounding East Pacific Ocean. However, no differences between male and female migration patterns were observed.

Comparisons of genetic markers with uni- and bi-parental inheritance are often used to determine the fidelity of females versus males to particular groupings or locations. Higher pairwise population F_{ST} values based on mitochondrial as opposed to nuclear markers are usually interpreted as philopatric behavior in females. In Chapter 7, female gene flow (based on mitochondrial genes COX1 and NADH5) was compared to female+male gene flow (based on 11 nuclear microsatellites) between the Atlantic and other ocean basins. Results were inconclusive due to wide confidence intervals of microsatellite F_{ST} values. Due to an absence of significant population genetic structure within the Indian and Pacific Oceans, I was also not able to test for the occurrence of sex-biased dispersal in these regions.

Conservation threats and solutions

The attraction of mobulids to plankton-rich habitats and their distribution in the upper water column makes them vulnerable to capture in a wide array of fishing gear, including

recreational and commercial harpoon, gill net, longline, trawl, purse seine and trap fisheries (Chapter 7). Since the 1990's, the market for mobulid filter plates used in Chinese medicine has grown with dried products fetching prices of up to US\$71 kg⁻¹. This has led to a rapid expansion of targeted mobulid fisheries, with dried filter plates being exported to mainland China from Sri Lanka, the Philippines, Indonesia and India, where annual catches reach into the tens of thousands (Croll *et al.* submitted). Additionally, bycatch of mobulids in small-scale fisheries has been reported from various countries with, for example, a total of 1,600 mobulids taken in Indonesian skipjack tuna fisheries and around 8,000 individuals in Peruvian artisanal fisheries (Croll *et al.* submitted). In large-scale tuna purse seine fisheries, an estimated total of 13,900 mobulids is caught annually worldwide.

Long-term demographic changes in long-lived pelagic species are difficult to detect (Lewison *et al.* 2004). Time lag, a result of long generation times, can delay the response of a population by many years. Additionally, general difficulties associated with surveys of pelagic animals make the detection of population trends extremely challenging. However, global harvest of mobulids is increasing, and at the same time, decreasing catch rates are reported from various regions. Indeed, population declines have already been reported from several countries, e.g. Mozambique, the Philippines, Indonesia, Mexico and India (Croll *et al.* submitted). As a result, sustainability of current mobulid catch-rates is unlikely.

The pelagic and highly vagile nature of mobulids also impedes conservation efforts. Mobulids roam widely, and may well spend their lives in different exclusive economic zones (and/or the open ocean), where they face a myriad of different threats from targeted and bycatch fisheries. Effective conservation will therefore require coordinated actions by fisheries stakeholders, international trade organizations, non-governmental conservation groups and consumer organizations. Based on the review of the main sources of mortality in mobulids (Chapter 7), three key recommendations have the potential to provide significant conservation benefits:

1. Reduce the demand for filter plates used in Chinese medicine through education and (if necessary) through boycotts and embargos;
2. Reduce bycatch in commercial tuna purse seine fisheries by enforcing fishing prohibitions in important regions such as feeding, breeding and pupping grounds, and by development of net designs that prevent bycatch and/or allow for effective live-release of captured animals;
3. Reduce targeted and bycatch of mobulids in artisanal fisheries through fisheries regulations and technical assistance with gear modification and improvement of live-release techniques.

Samenvatting

SAMENVATTING

Duivelsroggen (de collectieve naam voor *Manta* en *Mobula* roggen van de Familie Mobulidae) zijn grote pelagische planktoneters (filteraars) en behoren, door hun langzame voortplanting en extreem lage vruchtbaarheid, tot de meest kwetsbare haaiachtigen. Helaas bestaat er een intensieve en zich uitbreidende visserij in talloze Aziatische landen, die gedreven wordt door het gebruik van filterplaten in de Chinese Geneeskunde. Duivelsroggen worden ook frequent en over hun hele verspreidingsgebied gevangen als bijvangst in de tonijnvisserij. Door de lage intrinsieke groei van duivelsrogpopulaties kunnen zelfs lage vangsten tot significante negatieve populatietrends lijden.

De evolutionaire geschiedenis van duivelsroggen is lang onduidelijk gebleven, ondanks een recente interesse in de ecologie en bescherming van deze groep roggen. Deze thesis is ontworpen om een basis aan kennis te verwerven over duivelsrog evolutie en populatie genetica, zodat management- en beschermingsmaatregelen beter geïnformeerd en doeltreffender kunnen worden uitgevoerd. Ik heb Next Generation Sequencing (NGS) van hele mitochondriële genomen, conventionele Sanger sequencing en aanvullende GenBank sequenties gebruikt om, voor de eerste keer, de evolutionaire relaties tussen alle momenteel beschreven duivelsroggen te reconstrueren. Ook heb ik patronen in genetische variatie geanalyseerd binnen een van de meest frequent gevangen duivelsroggen, de wereldwijd verspreide gestekelde duivelsrog, *Mobula japanica*.

De belangrijkste vragen in deze thesis waren:

1. Wat zijn de evolutionaire relaties binnen de Family Mobulidae, en wat is de geologische timing en patroon van hun radiatie? Wat zijn waarschijnlijke mechanismen en de drijvende krachten achter Mobulidae soortvorming?
2. Wat zijn de geografische schaal en verbinding tussen genetisch gedefinieerde populaties van de gestekelde duivelsrog, *M. japanica*? Wat waren de invloeden van historische klimaatsverandering op de historische populatie structuur en demografie?
3. Is er bewijs voor verschillende verspreidingspatronen tussen de verschillende geslachten in *M. japanica*?
4. Wat zijn de belangrijkste antropogene dreigingen voor duivelsroggen? Hoe kan deze groep soorten beter beschermd worden?

Door de opkomst van Next Generation Sequencing (NGS) is het nu mogelijk om relatief goedkoop en efficiënt grotere datasets te genereren, in vergelijking met traditioneel Sanger sequenzen of microsatelliet analyse. De bijna onbegrensde mogelijkheden van NGS zorgen voor een ongekende resolutie binnen de biologische wetenschappen, onder andere in intra- en inter-specifieke studies. De toepassing in deze thesis van NGS-gegenereerde mitogenomen zorgde voor zowel sterk onderbouwde fylogenetische patronen, als ook de ontdekking van vier mitogenomische groepen binnen *M. japanica*, in tegenstelling met de toepassing van losse mitochondriële genen en microsatelliet markers.

Fylogenie van duivelsroggen

Ondanks de formele scheiding van duivelsroggen in twee genera (i.e. *Manta* en *Mobula*) heeft mijn onderzoek aangetoond dat het genus *Mobula* parafoyletisch is door de toevoeging van *Manta* (Hoofdstuk 5). Alhoewel parafoylie van duivelsroggen al eerder was voorgesteld op basis van morfologie (Adnet *et al.* 2012; Aschliman *et al.* 2012b; Gonzalez-Isais & Dominguez 2004; Herman *et al.* 2000), parasitaire evolutie (Benz & Deets 1988) en de vergelijkingen van DNA sequenties van een klein aantal duivelsroggen (Aschliman *et al.* 2012a; Naylor *et al.* 2012), is dit de eerste keer dat parafoylie bevestigd is door DNA sequenties van alle duivelsroggen. Mijn onderzoek laat verder zien dat duivelsroggen ingedeeld zijn in drie gemakkelijk te onderscheiden groepen, of 'clades' (Figuur 1). Op basis van een fossiel gecalibreerde fylogenetische analyse schat ik het tempo van de mitochondriële evolutie in duivelsroggen op 9.93×10^{-9} nucleotide substituties per jaar per site ($S s^{-1} j^{-1}$), wat ongeveer een orde van grootte langzamer is dan die in zoogdieren (Martin *et al.* 1992), maar overeenkomt met schattingen voor andere haai-achtigen (Dudgeon *et al.* 2012). Gebaseerd op dit tempo zijn duivelsroggen van de nauwverwante koeneusroggen gedivergeerd tijdens het Oligoceen, ongeveer 30 miljoen jaar geleden. Deze divergentie markeerde de transitie van een durofage (verpletteren van schelpen) naar een filtervoeding leefwijze. Daaropvolgende divergenties van soorten vonden plaats tijdens het Vroege and Midden Mioceen (19-17 miljoen jaar geleden), gevolgd door lange interne takken en een tweede interval van divergentie tijdens het Pliocene en Pleistoceen (3.6 Mya – recent) (Figure 1). Ik speculeer dat opwelling van diep nutriëntrijk water, wat direct de beschikbaarheid van eten beïnvloedt, een belangrijke rol in het ontstaan en divergeren van de duivelsroggen heeft gespeeld (zie hieronder).


Fylogeography en populatie genetische structuur van *Mobula japonica*

In Hoofdstuk 6 behandel ik de geografische schaal van genetisch geïsoleerde populaties, als ook de demografische geschiedenis van populaties van een van de 11 duivelsroggen, de globaal voorkomende *M. japonica*. Populatie genetische structuur werd geëvalueerd met behulp van 263 individuen uit een totaal aan acht locaties in the Stille, Indische en Atlantische oceanen, in combinatie met Sanger sequenties van mitochondriële genen COX1 en NADH5 en 11 microsatelliet loci. De resultaten suggereren hoge connectiviteit over uitgespreide oceanische gebieden. De Atlantische Oceaan is welliswaar genetisch gedifferentieerd van de Indische en Stille Oceanen (paarsgewijze $F_{ST}(\text{msats})=0.033-0.046$; paarsgewijze $F_{ST}(\text{mtDNA})=0.140-0.167$), maar binnen de Indische en Stille Oceanen lieten analyses zien dat er geen populatie structuur aanwezig is. Alleen een hiërarchische AMOVA analyse resulteerde in een zwakke genetische structuur ($F_{ST}(\text{msats})=0.007$, $F_{ST}(\text{mtDNA})=0.048$) tussen twee populaties uit de Oostelijke Stille Oceaan, en tussen de Atlantische en de Indische/Stille Oceanen. Er was geen bewijs voor isolatie door afstand (Isolation by Distance).

Een samenvatting van populatiestructuur in andere haaiachtigen suggereert dat de migratiecapaciteit van *M. japonica* alleen wordt overtroffen door die van de grootsten van de pelagische haaien, de walvishaai (Castro *et al.* 2007; Vignaud *et al.* 2014) en de reuzenhaai (Hoelzel *et al.* 2006). De gelijkenis in populatiestructuur tussen deze soorten en *M. japonica* was verassend, gezien dat zowel bij de walvishaai als de reuzenhaai transoceanische migraties voorkomen (Eckert *et al.* 2002; Gore *et al.* 2008). *M. japonica* daarentegen laat veel kortere seizoensgebonden migratie zien: satellietzender data suggereert migratie tussen de Golf van California en de aangrenzende Oostelijke Stille Oceaan, over een afstand van minder dan 1500 km. Discrepancies tussen direct waargenomen migratieafstanden en indirect geschatte afstanden gebaseerd op genetische data zijn gebruikelijk. Satellietzender studies volgen individuen gedurende maan-

den tot jaren, terwijl populatie genetische data verschillende generaties beschrijven (tientallen jaren). Het is mogelijk dat gedurende El Niño jaren, wanneer opwelling in de gebieden waar *M. japanica* frequent voorkomt zwaar gereduceerd is, de soort over veel langere afstanden zwemt naar gebieden met hogere productie, waardoor migratie overijd uit elkaar liggende regio's gepromoot wordt.

In een poging eventuele verborgen populatie structuur zichtbaar te maken zijn hele mitochondriële genomen gesequenced met behulp van NGS. Dit verhoogde de resolutie dramatisch (11,427 bp) in vergelijking tot de twee-gen mtDNA dataset (COX1/NADH5, 1107 bp). Alhoewel populatie genetische patronen in de mitogenomen overeenkwamen met die gebaseerd op de kleinere dataset, zijn *M. japanica* individuen verdeeld in vier groepen, waarbij groepslidmaatschap maar in kleine mate bepaald is door locatie van herkomst (Figuur 2) (Hoofdstuk 5). De vier groepen zijn waarschijnlijk gedivergeerd tijdens het Late Pleistoceen (126 –11.7 Kya), gebaseerd op sequentie divergentie en een mitochondriële klok van $9.93 \times 10^{-9} \text{ S s}^{-1} \text{ j}^{-1}$. Ik speculeer dat deze groepsstructuur het resultaat is van divergentie door fysieke migratie barrières die in het Pleistoceen aanwezig waren, zoals de koude wateren rond de Kaap de Goeie Hoop tijdens perioden van lage 'Agulhas leakage' (Peeters 2002), en de Sunda/Sahul landbrug tijdens de laatste ijstijd (Clark *et al.* 2009). Beide barrières zijn eerder al geïmpliceerd in de formatie van mitochondriële groepen in andere migrerende tropische pelagische soorten, bijvoorbeeld in tonijnen, zwaardvissen, cetaceën en in sommige van de grotere pelagische haaiensoorten (Amaral *et al.* 2012; Bremer *et al.* 1998; Daly-Engel *et al.* 2012; Duncan *et al.* 2006; Martinez *et al.* 2006; Morin *et al.* 2010; Rosel & Block 1996; Vinas *et al.* 2004). Het feit dat groepslidmaatschap maar losjes gebaseerd was op geografische locatie van herkomst impliceert periodes van




Figuur 1: Divergentietijd schattingen op basis van een analyse van mitogenomen. Puntschattingen van divergentietijden zijn aangegeven boven elk knooppunt, en transparante blauwe horizontale strepen indiceren tijd (95% highest posterior density). Kleuren van lijnen tussen taxa laten de relatieve snelheid van nucleotidesubstitutie zien (blauw is langzaam, paars is gemiddeld, rood is snel). Rode strepen onder de tijdperkenbalk illustreren perioden van klimaat warming and geassocieerde lage opwelling intensiteiten. De zwarte lijnen daaronder geven de timing van het voorkomen van uitgestorven en bestaande duivelsrogsoorten weer, gebaseerd op fossielen.

hoge connectiviteit tussen geïsoleerde regio's, waarbij de verschillende mitogenoomgroepen zich weer mengden. De vier groepen waren niet te onderscheiden van elkaar op basis van microsatelliet loci, wat impliceert dat individuen van verschillende groepen tot dezelfde soort behoren. Verder lieten drie van de vier groepen bewijs van demografische expansie zien, rond 40 duizend jaar geleden, en tussen 10-25 duizend jaar geleden. Perioden van toenemende opwelling en een daarmee samenhangende hoge primaire productie die onder invloed van het klimaatsveranderingen voorkwamen, zouden gezorgd kunnen hebben voor verhoogde draagkracht voor een pelagische planktoneter.

Geslachts-afhankelijke verspreiding

Veel haaiachtigen vertonen filopatrisch gedrag, wat inhoudt dat individuen terugkeren naar de plek van geboorte of een andere geadopteerde locatie om zich voort te planten (Mayr 1963). Zulk gedrag is bij haaiachtigen vaak geslachts-gebonden, waarbij vooral het vrouwelijke geslacht herhaaldelijk terugkeert naar specifieke locaties voor reproductie of om jongen te werpen, terwijl het mannelijke geslacht over veel grote gebieden migreert (bijvoorbeeld zwartpunthaaien, citroenhaaien en witte haaien, samengevat door Hueter *et al.* 2004).



Figuur 2: Bayesian fylogenetische boom gebaseerd op mitogenoomsequenties van 60 *Mobula japonica* individuen uit Mexico (Stille Oceaan, PAMX)), Costa Rica (Stille Oceaan, PACR), de Indische Oceaan (IO), de Oostelijke Atlantische Oceaan (EAO), een *Mobula mobular* individu (MMO), en een individu van elk van de twee 'outgroup' soorten (*Mobula tarapacana* en *M. birostris*). De kleur van ijenen correspondeert met locatie van herkomst. Posterior probabilities (PP) voor knooppunten die de groepen van elkaar scheiden staan vermeld; alle andere PP zijn hoger dan 0.80 (niet vermeld). De gele ster indiceert de positie van de outgroup soorten.

Filopatrisch gedrag heeft belangrijke implicaties voor het behoud van een soort, omdat voorplantings- en geboortegronden belangrijke habitats vormen voor een populatie, zoals ook geboorterivieren voor zalmen en broedstranden voor zeeschildpadden. Er bestaat geen betrouwbare informatie over het verschijnsel filopatrie in *M. japonica*, maar vangst- en satellitzenderdata suggereren dat seizoensgebonden migratie plaatsvindt tussen regio's van hoge productiviteit in de Golf van California en de aangrenzende Oostelijk Stille Oceaan. Er zijn echter geen verschillen tussen de geslachten geobserveerd.

Vergelijkingen tussen genetische markers met alleen vrouwelijke (mitochondrieel DNA), en met zowel mannelijke als vrouwelijke overerving (nucleair DNA) worden vaak gebruikt om te bepalen of een geslacht van een soort meer trouw is aan een bepaalde groep of locatie. Hogere paarsgewijze populatie F_{ST} waarden gebaseerd op mitochondriële versus nucleaire markers worden gewoonlijk geïnterpreteerd als filopatrisch gedrag in de vrouwelijke sekse. In Hoofdstuk 7 wordt de uitwisseling van vrouwelijke genen (gebaseerd op mitochondriële genen COX1 en NADH5) vergeleken met genen die door beide geslachten worden doorgegeven (de 11 nucleaire microsatellieten), tussen de Atlantische en andere Oceanen. Resultaten waren niet eenduidig door de wijde betrouwbaarheidsintervallen van de microsatelliet F_{ST} waarden. Door de afwezigheid van significante populatiestructuur binnen de Indische en Stille Oceanen was het ook niet mogelijk om de aanwezigheid van filopatrisch gedrag in deze regio's te onderzoeken.

Behoud van duivelsroggen: problemen en oplossingen

De aantrekkingskracht van planktonrijke gebieden op duivelsroggen, samen met hun distributie in de bovenste waterkolom, maken deze soorten uiterst kwetsbaar voor een groot aantal verschillende visserijen, onder andere recreatieve en commerciële harpoenvisserij, kieuw-netvisserij, langlijnvisserij, trawlvisserij, ringzegenvisserij en kooivisserij (Hoofdstuk 7). De markt voor filterplaten van duivelsroggen, welke gebruikt worden in de Chinese Geneeskunde, is sterk gegroeid sinds 1990, en prijzen voor gedroogde producten liggen nu rond de €65 per kilo. Dit heeft geleid tot een snelle expansie van een gerichte duivelsrogvisserij in Sri Lanka, de Filipijnen, Indonesië en India, vanwaar gedroogde filterplaten worden geëxporteerd naar het Chinese vasteland (Croll *et al.* Submitted). In de ringzegenvisserij naar tonijn worden ook nog eens gemiddeld 13,900 duivelsroggen per jaar gevangen.

Het is moeilijk om langdurige veranderingen in de demografie van langlevende soorten te detecteren. Als resultaat van de lange generatietijd kan verhoogde mortaliteit pas na vele jaren een effect geven. Ook is het, door de enorme uitgestrektheid en ontoegankelijkheid van de pelagische habitat over het algemeen moeilijk om populatietrends in zulke soorten te onderzoeken. Toch is het duidelijk dat de globale oogst van duivelsroggen toeneemt, terwijl tegelijkertijd de vangstratio's in uiteenlopende gebieden omlaag gaan. Negatieve populatietrends zijn daadwerkelijk waargenomen in verschillende landen, onder andere Mozambique, de Filipijnen, Indonesië, Mexico en India (Croll *et al.* Submitted). Het is daarom onwaarschijnlijk dat de huidige vangst van duivelsroggen duurzaam is.

De pelagische en de vrij-zwemmende natuur van duivelsroggen zorgt voor meer problemen omtrent het behoud van deze groep. Duivelsroggen hebben een wijd verspreidingsgebied, en het is daarom mogelijk dat ze hun leven binnen verschillende exclusieve economische zones (en/of de open oceaan) doorbrengen, waar ze potentieel in aanraking komen met bedreigingen van gerichte- of bijvangst visserijen. Effectief behoud van deze groep vraagt daarom gecoördineerde acties van visserijaandeelhouders, internationale handelsorganisaties, natuurbehoud groepen en consumentenorganisaties. Gebaseerd op een samenvatting van de belangrijkste sterfteoorzaken in duivelsroggen (Chapter 7) kunnen drie aanbevelingen gegeven worden die potentieel het behoud van deze soorten ten goede komen:

1. Reduceer de vraag naar filterplaten vanuit Chinese Geneeskunde, door educatie en (zo nodig) door boycotts en embargo's;
2. Reduceer bijvangst in de commerciële ringzegenvisserij naar tonijnen, door het afdwingen van een verbod op visserij in belangrijke regio's, zoals foerageer-, voortplantings- en geboortegronden, en door netten te ontwikkelen die bijvangst voorkomen en/of toestaan dat dieren op een effectieve manier levend vrijgelaten kunnen worden.
3. Reduceer gerichte- en bijvangst van duivelsroggen in kleinschalige visserij door regelgeving, en door technische assistentie te verlenen met de modificatie van vistuig en het verbeteren van technieken om dieren levend vrij te laten.

References

A

- Acebes JMV (2009) Historical whaling in the Philippines: origins of 'indigenous subsistence whaling', mapping whaling grounds and comparison with current known distribution: a HMAP Asia Project Paper.
- Adimey NM, Hudak CA, Powell JR, Bassos-Hull K, Foley A, Farmer NA, White L, Minch K (2014) Fishery gear interactions from stranded bottlenose dolphins, Florida manatees and sea turtles in Florida, U.S.A. *Marine Pollution Bulletin* **81**, 103-115.
- Adnet S, Cappetta H, Guinot G, Notarbartolo di Scara G (2012) Evolutionary history of the devil-rays (Chondrichthyes: Myliobatiformes) from fossil and morphological inference. *Zoological Journal of the Linnean Society* **166**, 132-159.
- Ahonen H, Harcourt RG, Stow AJ (2009) Nuclear and mitochondrial DNA reveals isolation of imperilled grey nurse shark populations (*Carcharias taurus*). *Molecular Ecology* **18**, 4409-4421.
- Alava MNR, Dolumbalo ERZ, Yaptinchay AA, Trono RB (2002) Fishery and trade of whale sharks and manta rays in the Bohol Sea, Philippines. In: Elasmobranch biodiversity, conservation, and management: Proceedings of the international seminar and workshop, Sabah, Malaysia, July 1997 eds. Fowler SL, Reed TM, Dipper FA), pp. 132-148. IUCN SSC Shark Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.
- Alexander A, Steel D, Slikas B, Hoekzema K, Carraher C, Parks M, Cronn R, Baker CS (2013) Low diversity in the mitogenome of sperm whales revealed by Next-Generation Sequencing. *Genome Biology and Evolution* **5**, 113-129.
- Alongi DM, Wirasantosa S, Wagey T, Trott LA (2012) Early diagenetic processes in relation to river discharge and coastal upwelling in the Aru Sea, Indonesia. *Marine Chemistry* **140-141**, 10-23.
- Alvarado Bremer JR, Vinas J, Mejuto J, Ely B, Pla C (2005) Comparative phylogeography of Atlantic bluefin tuna and swordfish: the combined effects of vicariance, secondary contact, introgression, and population expansion on the regional phylogenies of two highly migratory pelagic fishes. *Molecular Phylogenetics Evolution* **36**, 169-187.
- Amandè J, Ariz J, Chassot E (2008) Bycatch of the European purse seine tuna fishery in the Atlantic Ocean for the 2003–2007 period. Report IOTC-2008-WPEB-12. IOTC Ecosystem and By-catch Working Group, Bangkok, Thailand.
- Amandè J, Ariz J, Chassot E, Chavance P, Molina ADD, Gaertner D, Murua H, Pianet R, Ruiz J (2011) By-catch and discards of the European purse seine tuna fishery in the Atlantic Ocean: estimation and characteristics for 2008 and 2009. *Collective Volume Scientific Papers ICCAT* **66**, 2113-2120.
- Amandè MJ, Ariz J, Chassot E, de Molina AD, Gaertner D, Murua H, Pianet R, Ruiz J, Chavance P (2010) Bycatch of the European purse seine tuna fishery in the Atlantic Ocean for the 2003–2007 period. *Aquatic Living Resources* **23**, 353-362.
- Amaral AR, Beheregaray LB, Bilgmann K, Freitas L, Robertson KM, Sequeira M, Stockin KA, Coelho MM, Moller LM (2012) Influences of past climatic changes on historical population structure and demography of a cosmopolitan marine predator, the common dolphin (genus *Delphinus*). *Molecular Ecology* **21**, 4854-4871.
- Anderson RC, Adam MS, Goes JI (2011) From monsoons to mantas: seasonal distribution of *Manta alfredi* in the Maldives. *Fisheries Oceanography* **20**, 104-113.
- Anderson RF, Ali S, Bradtmiller LI, Nielsen SHH, Fleisher MQ, Anderson BE, Burckle LH (2009) Wind-driven upwelling in the Southern Ocean and the deglacial rise in atmospheric CO₂. *Science* **323**, 1443-1448.
- Andreasen DJ, Ravelo AC (1997) Tropical Pacific Ocean thermocline depth reconstructions for the Last Glacial Maximum. *Paleoceanography* **12**, 395-413.

- Arlyza IS, Shen KN, Durand JD, Borsa P (2013) Mitochondrial haplotypes indicate parapatric-like phylogeographic structure in blue-spotted maskray (*Neotrygon kuhlii*) from the Coral Triangle Region. *Journal of Heredity*, 725-733.
- Aschliman NC (2014) Interrelationships of the durophagous stingrays (Batoidea: Myliobatidae). *Environmental Biology of Fishes* **97**, 967-979.
- Aschliman NC, Claeson KM, McEachran JD (2012a) Phylogeny of Batoidea. In: *Biology of sharks and their relatives* (eds. Carrier JC, A. MJ, R. HM). CRC Press, Boca Raton.
- Aschliman NC, Nishida M, Miya M, Inoue JG, Rosana KM, Naylor GJ (2012b) Body plan convergence in the evolution of skates and rays (Chondrichthyes: Batoidea). *Molecular Phylogenetics and Evolution* **63**, 28-42.
- Avice JC (1989) Gene trees and organismal histories: a phylogenetic approach to population biology. *Evolution* **43**, 1192-1208.
- Avice JC (2000) *Phylogeography. The history and formation of species* Harvard University Press, Cambridge, MA.
- Avice JC (2008) The history, purview, and future of conservation genetics. In: *Conservation biology: evolution in action* (eds. Carroll SP, Fox C), pp. 5-15. Oxford University Press, New York.
- Avice JC (2010) Perspective: conservation genetics enters the genomics era. *Conservation Genetics* **11**, 665-669.
- Ayala L, Amoros S, Cespedes C (2008) Catch and by-catch of albatross and petrel in longline and gillnet fisheries in Northern Peru. Final report to the Rufford Small Grants for Nature Conservation. APECO, Lima, Peru.

B

- Baker CS, Palumbi SR (1994) Which whales are hunted? A molecular genetic approach to monitoring whaling. *Science* **265**, 1538-1539.
- Bakun A (1990) Global climate change and intensification of coastal ocean upwelling. *Science* **247**, 198-201.
- Bancroft EN (1831) On several fishes of Jamaica. *Proceedings of the Zoological Society of London* **1**, 134.
- Barrett RDH, Schluter D (2008) Adaptation from standing genetic variation. *Trends in Ecology and Evolution* **23**, 38-44.
- Becquet C, Przeworski M (2009) Learning about modes of speciation by computational approaches. *Evolution* **63**, 2547-2562.
- Benton MJ, Asher RJ (2009) Calibrating and constraining molecular clocks. In: *Timetree of life* (eds. Hedges SB, Kumar S), pp. 35-86. Oxford University Press, Oxford.
- Benz GW, Deets GB (1988) Fifty-one years later: an update on *Entepherus*, with a phylogenetic analysis of *Cecropidae* Dana, 1849 (Copepoda: Siphonostomatoida). *Canadian Journal of Fisheries and Aquatic Sciences* **66**, 856-865.
- Berman-Kowalewski M, Gulland FMD, Wilkin S, Calambokidis J, Mate B, Cordaro J, Rotstein D, Leger JS, Collins P, Fahy K, Dover S (2010) Association between blue whale (*Balaenoptera musculus*) mortality and ship strikes along the California coast. *Aquatic Mammals* **36**, 59-66.
- Bernardi G, Holbrook SJ, Schmitt RJ (2001) Gene flow at three spatial scales in a coral reef fish, the three-spot dascyllus, *Dascyllus trimaculatus*. *Marine Biology* **138**, 457-465.
- Bigelow HB, Schroeder WC (1953) Sawfishes, guitarfishes, skates, and rays. In: *Fishes of the western North Atlantic* (eds. Bigelow HB, Schroeder WC), pp. 508-514. Sears Foundation for Marine Research, Yale University, New Haven, CT.
- Birky CW, Fuerst P, Maruyama T (1989) Organelle gene diversity under migration, mutation, and drift - equilibrium expectations, approach to equilibrium, effects of heteroplasmic cells,

- and comparison to nuclear genes. *Genetics* **121**, 613-627.
- Bizzarro J (2001) The occurrence of mobulid (Chondrichthyes: Mobulidae) rays in the Gulf of California and their associated fisheries. In: Technical report prepared for Monterey Bay Aquarium. Monterey Bay Aquarium, Monterey, U.S.A.
- Bizzarro JJ, Smith WD, Hueter RE, Tyminski J, Márquez-Farías JF, Castillo-Géniz JL, Cailliet GM, Villavicencio-Garayzar CJ (2007) The status of shark and ray fishery resources in the Gulf of California: applied research to improve management and conservation. In: Technical Report 2009-01. Moss Landing Marine Laboratories, Moss Landing, U.S.A.
- Blakey RC (2008) Gondwana paleogeography from assembly to breakup - A 500 m.y. odyssey. *The Geological Society of America* **441**, 1-28.
- Bleeker P (1859) Enumeratio specierum piscium hucusque in Archipelago Indico observatarum. *Acta Societatis Scientiarum Indiarum Neerlandensium* **3**, 1-6.
- Blower DC, Pandolfi JM, Bruce BD, Gomez-Cabrera M, Ovenden JR (2012) Population genetics of Australian white sharks reveals fine-scale spatial structure, transoceanic dispersal events and low effective population sizes. *Marine Ecology Progress Series* **455**, 229-244.
- Boersma A, Silva IP (1991) Distribution of Paleogene planktonic foraminifera - analogies with the Recent? *Palaeogeography, Palaeoclimatology, Palaeoecology* **83**, 29-48.
- Bolliet T, Holbourn A, Kuhnt W, Laj C, Kissel C, Beaufort L, Kienast M, Andersen N, Garbe-Schonberg D (2011) Mindanao Dome variability over the last 160 kyr: episodic glacial cooling of the West Pacific Warm Pool. *Paleoceanography* **26**, PA1208.
- Bonnaterre JP (1788) *Tableau encyclopédique et méthodique des trois regnes de la nature* Chez Panckoucke, Paris.
- Boore JL (1999) Animal mitochondrial genomes. *Nucleic Acids Research* **27**, 1767-1780.
- Bourdon J (1999) A fossil Manta from the Early Pliocene (Zanclean) of North America. *Tertiary Research* **19**, 79-84.
- Bourjea J, Lapegue S, Gagnevin L, Broderick D, Mortimer JA, Ciccione S, Roos D, Taquet C, Grizel H (2007) Phylogeography of the green turtle, *Chelonia mydas*, in the southwest Indian Ocean. *Molecular Ecology Notes* **16**, 175-186.
- Bowen BW, Bass AL, Soares L, Toonen RJ (2005) Conservation implications of complex population structure: lessons from the loggerhead turtle (*Caretta caretta*). *Mol Ecol* **14**, 2389-2402.
- Bowen BW, Karl SA (2007) Population genetics and phylogeography of sea turtles. *Molecular Ecology* **16**, 4886-4907.
- Bremer JRA, Stequert B, Robertson NW, Ely B (1998) Genetic evidence for inter-oceanic subdivision of bigeye tuna (*Thunnus obesus*) populations. *Marine Biology* **132**, 547-557.
- Britten RJ (1986) Rates of DNA sequence evolution differ between taxonomic groups. *Science* **231**, 1393-1398.
- Brown WM, Prager EM, Wang A, Wilson AC (1982) Mitochondrial DNA sequences of primates: tempo and mode of evolution. *Journal of Molecular Evolution* **18**, 225-239.
- Buroker NE, Brown JR, Gilbert TA, O'Hara PJ, Beckenbach AT, Thomase WK, Smith MJ (1990) Length heteroplasmy of sturgeon mitochondrial DNA: an illegitimate elongation model. *Genetics* **124**, 157-163.
- Bustamante C, Couturier LIE, Bennett MB (2012) First record of *Mobula japanica* (Rajiformes: Myliobatidae) from the south-eastern Pacific Ocean. *Marine Biodiversity Records* **5**, e48.

C

- Camhi M, Valenti S, Fordham S, Fowler S, Gibson C (2009) The conservation status of pelagic sharks and rays: report of the IUCN shark specialist group pelagic shark red list workshop. In: IUCN Species Survival Commission's Shark Specialist Group.

- Canese S, Cardinali A, Romeo T, Giusti M, Salvati E, Angiolillo M, Greco S (2011) Diving behavior of the giant devil ray in the Mediterranean Sea. *Endangered Species Research* **14**, 171-176.
- Cappetta H (1975) Contribution à l'étude des Sélaciens du groupe Monmouth (Campanien-Maastrichtien) du New Jersey *Palaeontographica, Abteilung A* **151**, 1-46.
- Cappetta H (1985) Sur une nouvelle espèce de *Burnhamia* (Batomorphii, Mobulidae) dans l'Yprésien des Ouled Abdoun, Maroc. *Tertiary Research* **7**, 27-33.
- Cappetta H, Stringer GL (1970) A new batoid genus (Neoselachii: Myliobatiformes) from the Yazoo Clay (Upper Eocene) of Louisiana, U.S.A. *Tertiary Research* **21**, 51-56.
- Cardeñoso D, Hyde J, Caballero S (2014) Genetic diversity and population structure of the pelagic thresher shark (*Alopias pelagicus*) in the Pacific Ocean: evidence for two Evolutionarily Significant Units. *PLoS One* **9**, e110193.
- Carvalho GR, Hauser L (1994) Molecular genetics and the stock concept in fisheries. *Reviews in Fish Biology and Fisheries* **4**, 326-350.
- Case GR (1980) A selachian fauna from the Trent Formation, Lower Miocene (Aquitania) of Eastern North Carolina. *Palaeontographica, Abteilung A* **171**, 75-103.
- Castillo-Paez A, Sosa-Nishizaki O, Galvan-Magana F, Blanco-Parra MD, Rocha-Olivares A (2014) Strong population structure and shallow mitochondrial phylogeny in the banded guitarfish, *Zapteryx exasperata* (Jordan y Gilbert, 1880), from the Northern Mexican Pacific. *Journal of Heredity* **105**, 91-100.
- Castro AL, Stewart BS, Wilson SG, Hueter RE, Meekan MG, Motta PJ, Bowen BW, Karl SA (2007) Population genetic structure of Earth's largest fish, the whale shark (*Rhincodon typus*). *Molecular Ecology* **16**, 5183-5192.
- Chabot CL, Allen LG (2009) Global population structure of the tope (*Galeorhinus galeus*) inferred by mitochondrial Control Region sequence data. *Molecular Ecology* **18**, 545-552.
- Chappell J, Shackleton NJ (1986) Oxygen isotopes and sea level. *Nature* **324**, 137-140.
- Charif D, Lobry J (2007) SeqinR 1.0-2: a contributed package to the R project for statistical computing devoted to biological sequences retrieval and analysis. In: *Structural approaches to sequence evolution: molecules, networks, populations* (eds. Bastolla U, Porto M, Roman H, Vendruscolo M), pp. 207-232. Springer Verlag, New York, USA.
- Charlesworth J, Eyre-Walker A (2007) The other side of the nearly neutral theory, evidence of slightly advantageous back-mutations. *Proceedings of the National Academy of Sciences U.S.A* **104**, 16992-16997.
- Chassot E, Dewals P, Floch L, Lucas V, Morales-Vargas M, Kaplan D (2010) Analysis of the effects of Somali piracy on the European tuna purse seine fisheries of the Indian Ocean. In: *Working document for IOTC. Indian Ocean Tuna Commission*.
- Chen C, Liu K, Joung S, Phipps MJ (2002) Taiwan's shark fishery - an overview. In: *Elasmobranch biodiversity, conservation, and management: Proceedings of the International Seminar and Workshop, Sabah, Malaysia, July 1997* eds. Fowler SL, Reed TM, Dipper FA), pp. 95-103. IUCN SSC Shark Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.
- Chevolot M, Hoarau G, Rijnsdorp AD, Stam WT, Olsen JL (2006) Phylogeography and population structure of thornback rays (*Raja clavata* L., Rajidae). *Molecular Ecology* **15**, 3693-3705.
- Chevolot M, Wolfs PHJ, Pálsson J, Rijnsdorp AD, Stam WT, Olsen JL (2007) Population structure and historical demography of the thorny skate (*Amblyraja radiata*, Rajidae) in the North Atlantic. *Marine Biology* **151**, 1275-1286.
- Chow S, Okamoto H, Miyabe N, Hiramatsu K, Barut N (2000) Genetic divergence between Atlantic and Indo-Pacific stocks of bigeye tuna (*Thunnus obesus*) and admixture around South Africa. *Molecular Ecology* **9**, 221-227.
- Cicimurri DJ, Knight JL (2009) Late Oligocene sharks and rays from the Chandler Bridge Formation, Dorchester County, South Carolina. *Acta Palaeontologica Polonica* **54**, 627-647.

- Claeson KM, O'Leary MA, Roberts EM, Sissoko F, Bouaré M, Tapanila L, Goodwin D, Gottfried MD (2010) First Mesozoic record of the stingray *Myliobatis wurnoensis* from Mali and a phylogenetic analysis of Myliobatidae incorporating dental characters. *Acta Palaeontologica Polonica* **55**, 655-674.
- Clark PU, Dyke AS, Shakun JD, Carlson AE, Clark J, Wohlfarth B, Mitrovica JX, Hostetler SW, McCabe AM (2009) The Last Glacial Maximum. *Science* **325**, 710-714.
- Clark T (2010) Abundance, home range, and movement patterns of manta rays (*Manta alfredi*, *M. birostris*) in Hawai'i, p. 150. University of Hawai'i, Manoa.
- Clarke SC, Magnussen J, Abercrombie DL, McAllister MK, Shivji MS (2006) Identification of Shark Species Composition and Proportion in the Hong Kong Shark Fin Market Based on Molecular Genetics and Trade Records. *Conservation Biology* **20**, 201-211.
- Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. *Molecular Ecology* **9**, 1657-1659.
- CMS (2014) Conservation of migratory sharks and rays. In: UNEP/CMS/Resolution 11.20, 11th Meeting of the Conference of the Parties, Quito, Ecuador.
- Coates AG, Aubry MP, Berggren WA, Collins LS, Kunk M (2003) Early Neogene history of the Central American arc from Bocas del Toro, western Panama. *Geological Society of America Bulletin* **115**, 271-287.
- Coates AG, Obando JA (1996) The geological evolution of the Central American Isthmus Chicago Press, Chicago.
- Coles RJ (1916) Natural history notes on the devil-fish, *Manta birostris* (Walbaum) and *Mobula olfersi* (Müller). *Bulletin American Museum of Natural History* **XXXV**, 649-657.
- Combes HJD, Caulet JP, Tribouillard N (2005) Monitoring the variations of the Socotra upwelling system during the last 250 kyr: A biogenic and geochemical approach. *Palaeogeography, Palaeoclimatology, Palaeoecology* **223**, 243-259.
- Compagno LJ (1999) An overview of Chondrichthyan systematics and biodiversity in Southern Africa. *Transactions of the Royal Society of South Africa* **54**, 75-120.
- Compagno LJ (2005) Checklist of living Chondrichthyes. In: Reproductive biology and phylogeny of Chondrichthyes. Sharks, batoids and chimaeras (ed. Hamlet WC), pp. 503-548. Science Publishers, Inc., Enfield.
- Compagno LJ, Last PR (1999a) Mobulidae. In: The living marine resources of the Western Central Pacific, pp. 1524-1529. FAO, Rome.
- Compagno LJ, Last PR (1999b) Mobulidae: devil rays. In: FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 3. Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae) eds. Carpenter KE, Niem VH), pp. 1524-1529, Rome, Italy.
- Cortés E, Papastamatiou YP, Carlson JK, Ferry-Graham L, Wetherbee BM, Cyrino JEP, Bureau DP, Kapoor BG (2008) An overview of the feeding ecology and physiology of elasmobranch fishes. In: Feeding and digestive functions of fishes (eds. Cyrino JEP, Bureau D, Kapoor BG), pp. 393-443. Science Publishers, Edenbridge.
- Couturier LI, Marshall AD, Jaine FR, Kashiwagi T, Pierce SJ, Townsend KA, Weeks SJ, Bennett MB, Richardson AJ (2012) Biology, ecology and conservation of the Mobulidae. *Journal of Fish Biology* **80**, 1075-1119.
- Couturier LIE, Bennett MB, Richardson AJ (2013) Mystery of giant rays off the Gaza strip solved. *Oryx* **47**, 480.
- Couturier LIE, Dudgeon CL, Pollock KH, Jaine FRA, Bennett MB, Townsend KA, Weeks SJ, Richardson AJ (2014) Population dynamics of the reef manta ray *Manta alfredi* in eastern Australia. *Coral Reefs* **33**, 329-342.
- Crandall ED, Frey MA, Grosberg RK, Barber PH (2008) Contrasting demographic history and phy-

- logeographical patterns in two Indo-Pacific gastropods. *Molecular Ecology* **17**, 611-626.
- Crisp MD, Cook LG (2009) Explosive radiation or cryptic mass extinction? Interpreting signatures in molecular phylogenies. *Evolution* **63**, 2257-2265.
- Croll DA, Dewar H, Dulvy NK, Fernando D, Malcolm F, Magana FG, Hall MA, Heinrichs S, Marshall A, McCauley D, Newton KM, Notarbartolo di Scara G, O'Malley M, O'Sullivan J, Poortvliet M, Roman M, Stevens G, Tershy BR, White WT (submitted) Slow life histories and fisheries impacts: The uncertain future of manta and devil rays. *Aquatic Conservation*.
- Croll DA, Marinovic B, Benson S, Chavez FP, Black N, Ternuilo R, Tershy B (2005) From wind to whales: trophic links in a coastal upwelling system. *Marine Ecology Progress Series* **289**, 117-130.
- Croll DA, Newton KM, Weng K, Galván-Magaña F, O'Sullivan J, Dewar H (2012) Movement and habitat use by the spine-tail devil ray in the Eastern Pacific Ocean. *Marine Ecology Progress Series* **465**, 193-200.
- Cuevas-Zimbrón E, Sosa-Nishizaki O, Pérez-Jiménez JC, O'Sullivan JB (2012) An analysis of the feasibility of using caudal vertebrae for ageing the spinetail devilray, *Mobula japanica* (Müller and Henle, 1841). *Environmental Biology of Fishes* **96**, 907-914.
- Cummings MP, Otto S, Wakely J (1995) Sampling properties of DNA sequence data in phylogenetic analysis. *Molecular Biology and Evolution* **12**, 814-822.
- Cunha RL, Coscia I, Madeira C, Mariani S, Stefanni S, Castilho R (2012) Ancient divergence in the trans-oceanic deep-sea shark *Centroscymnus crepidater*. *PLoS One* **7**, e49196.
- Curole JPFAUK, Kocher TD (1999) Mitogenomics: digging deeper with complete mitochondrial genomes. *Trends in Ecology and Evolution* **14**, 394-398.

D

- Daly-Engel TS, Seraphin KD, Holland KN, Coffey JP, Nance HA, Toonen RJ, Bowen BW (2012) Global phylogeography with mixed-marker analysis reveals male-mediated dispersal in the endangered scalloped hammerhead shark (*Sphyrna lewini*). *PLoS One* **7**, e29986.
- Darbyshire M (1967) The surface waters off the coast of Kerala, southwest India. *Deep-Sea Research* **14**, 295-320.
- de Carvalho MR, Maissey JG, Grande L (2004) Freshwater stingrays of the Green River formation of Wyoming (early Eocene), with the description of a new genus and species and an analysis of its phylogenetic relationships (Chondrichthyes: Myliobatiformes). *Bulletin American Museum of Natural History* **284**, 1-36.
- de Queiroz K, Gauthier J (1990) Phylogeny as a central principle in taxonomy: phylogenetic definitions of taxon names. *Systematic Zoology* **39**, 307-322.
- de Silva-Davila R, Palomares-Garcia JR (1998) Unusual larval growth production of *Nyctiphanes simplex* in Bahia de la Paz, Baja California Sur, Mexico. *Journal of Crustacean Biology* **18**, 490-498.
- Deakos MH, Baker JD, Bejder L (2011) Characteristics of a manta ray *Manta alfredi* population off Maui, Hawaii, and implications for management. *Marine Ecology Progress Series* **429**, 245-260.
- Defant A (1936) Das Kaltwasserauftriebsgebiet vor der Küste Südwestafrikas. *Landeskundliche Forschungen München Festschriften*, 52-66.
- Dewar H (2002) Preliminary Report : Manta Harvest in Lamakera. *Pfleger Institute of Environmental Research*, Oceanside, California.
- Dewar H, Mous P, Domeier M, Muljadi A, Pet J, Whitty J (2008) Movements and site fidelity of the giant manta ray, *Manta birostris*, in the Komodo Marine Park, Indonesia. *Marine Biology* **155**, 121-133.
- DeWoudy JA, Avise JC (2000) Microsatellite variation in marine, freshwater and anadromous

- fishes compared with other animals. *Journal of Fish Biology* **56**, 461-473.
- Diaz-Jaimes P, Uribe-Alcocer M, Rocha-Olivares A, Garcia-de-Leon FJ, Nortmoon P, Durand JD (2010) Global phylogeography of the dolphinfish (*Coryphaena hippurus*): The influence of large effective population size and recent dispersal on the divergence of a marine pelagic cosmopolitan species. *Molecular Phylogenetics and Evolution* **57**, 1209-1218.
- Diester-Haass L, Meyers PA, Laurence V (2002) The late Miocene onset of high productivity in the Benguela Current upwelling system as part of a global pattern. *Marine Geology* **180**, 87-103.
- Dixon MT, Hillis DM (1993) Ribosomal RNA secondary structure: compensatory mutations and implications for phylogenetic analysis. *Molecular Biology and Evolution* **10**, 256-267.
- Doherty PJ, Planes S, Mather P (1995) Gene flow and larval duration in 7 species of fish from the Great-Barrier-Reef. *Ecology* **76**, 2373-2391.
- Drummond AJ, Nicholls GK, Heled J, Rodrigo AG, Rodrigo W, Solomon W (2002) Estimating mutation parameters, population history and genealogy simultaneously from temporally spaced sequence data. *Genetics* **161**, 1307-1320.
- Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* **7**, 214.
- Drummond AJ, Rambaut A, Shapiro B, Pybus OG (2005) Bayesian coalescent inference of past population dynamics from molecular sequences. *Molecular Biology and Evolution* **22**, 1185-1192.
- Drummond AJ, Suchard MA (2010) Bayesian random local clocks, or one rate to rule them all. *BMC Biology* **8**, 114.
- Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* **29**, 1969-1973.
- Dubois N, Kienast M, Kienast S, Normandeau C, Calvert SE, Herbert TD, Mix A (2011) Millennial-scale variations in hydrography and biogeochemistry in the Eastern Equatorial Pacific over the last 100 kyr. *Quaternary Science Reviews* **30**, 210-223.
- Dudgeon CL, Blower DC, Broderick D, Giles JL, Holmes BJ, Kashiwagi T, Kruck NC, Morgan JA, Tillett BJ, Ovenden JR (2012) A review of the application of molecular genetics for fisheries management and conservation of sharks and rays. *Journal of Fish Biology* **80**, 1789-1843.
- Dudgeon CL, Broderick D, Ovenden JR (2009) IUCN classification zones concord with, but underestimate, the population genetic structure of the zebra shark *Stegostoma fasciatum* in the Indo-West Pacific. *Molecular Ecology* **18**, 248-261.
- Dudley SFJ, Cliff G (1993) Some effects of shark nets in the Natal nearshore environment. *Environmental Biology of Fishes* **36**, 243-255.
- Duffy CAJ, Abbott D (2003) Sightings of mobulid rays from northern New Zealand, with confirmation of the occurrence of *Manta birostris* in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* **37**, 715-721.
- Dulvy NK, Baum JK, Clarke S, Compagno LJV, Cortés E, Domingo A, Fordham S, Fowler S, Francis MP, Gibson C, Martínez J, Musick JA, Soldo A, Stevens JD, Valenti S (2008) You can swim but you can't hide: the global status and conservation of oceanic pelagic sharks and rays. *Aquatic Conservation: Marine and Freshwater Ecosystems* **18**, 459-482.
- Dulvy NK, Fowler SL, Musick JA, Cavanagh RD, Kyne PM, Harrison LR, Carlson JK, Davidson LN, Fordham SV, Francis MP, Pollock CM, Simpfendorfer CA, Burgess GH, Carpenter KE, Compagno LJ, Ebert DA, Gibson C, Heupel MR, Livingstone SR, Sanciangco JC, Stevens JD, Valenti S, White WT (2014a) Extinction risk and conservation of the world's sharks and rays. *eLife* **3**: e00590.
- Dulvy NK, Pardo SA, Simpfendorfer CA, Carlson JK (2014b) Diagnosing the dangerous demography of manta rays using life history theory. *PeerJ* **2**, e400.

- Duncan KM, Martin AP, Bowen BW, HG DEC (2006) Global phylogeography of the scalloped hammerhead shark (*Sphyrna lewini*). *Molecular Ecology* **15**, 2239-2251.
- Dunn KA, McEachran JD, Honeycutt RL (2003) Molecular phylogenetics of myliobatiform fishes (Chondrichthyes: Myliobatiformes), with comments on the effects of missing data on parsimony and likelihood. *Molecular Phylogenetics and Evolution* **27**, 259-270.
- Durrett R, Schweinsberg J (2004) Approximating selective sweeps. *Theoretical Population Biology* **66**, 129-138.

E

- Eckert SA, Dolar LL, Kooyman GL, Perrin W, Rahman RA (2002) Movements of whale sharks (*Rhincodon typus*) in South-east Asian waters as determined by satellite telemetry. *Journal of Zoology* **257**, 111-115.
- Edelaar P, Siepielski AM, Clobert J (2008) Matching habitat choice causes directed gene flow: a neglected dimension in evolution and ecology. *Evolution* **62**, 2462-2472.
- Ellegren H (2000) Microsatellite mutations in the germline: implications for evolutionary inference. *Trends in Genetics* **16**, 551-558.
- Elliott W (1846) *Carolina sports: by land and water: including incidents of devil-fishing*.
- Ely B, Vinas J, Bremer JRA, Black D, Lucas L, Covello K, Labrie AV, Thelen E (2005) Consequences of the historical demography on the global population structure of two highly migratory cosmopolitan marine fishes: the yellowfin tuna (*Thunnus albacares*) and the skipjack tuna (*Katsuwonus pelamis*). *BMC Evolutionary Biology* **5**.
- Escribano R, Daneri G, Farias L, Gallardo VA, Gonzalez HE, Gutierrez D, Lange CB, Morales CE, Pizarro O, Ulloa O, Braun M (2004) Biological and chemical consequences of the 1997-1998 El Niño in the Chilean coastal upwelling system: a synthesis. *Deep-Sea Research Part II: Topical Studies in Oceanography* **51**, 2389-2411.
- Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* **10**, 564-567.

F

- Faber JE, Stepien CA (1998) Tandemly Repeated Sequences in the Mitochondrial DNA Control Region and phylogeography of the pike-perches *Stizostedion*. *Molecular Phylogenetics and Evolution* **10**, 310-322.
- Fang G, Wang G, Fang Y, Fang W (2012) A review on the South China Sea western boundary current. *Acta Oceanologica Sinica* **31**, 1-10.
- Fedorov AV, Brierley CM, Lawrence KT, Liu Z, Dekens PS, Ravelo AC (2013) Patterns and mechanisms of early Pliocene warmth. *Nature* **496**, 43-49.
- Feldheim KA, Gruber SH, Ashley MV (2004) Reconstruction of parental microsatellite genotypes reveals female polyandry and philopatry in the lemon shark, *Negaprion brevirostris*. *Evolution* **58**, 2332-2342.
- Feldheim KA, Gruber SH, Dibattista JD, Babcock EA, Kessel ST, Hendry AP, Pikitch EK, Ashley MV, Chapman DD (2014) Two decades of genetic profiling yields first evidence of natal philopatry and long-term fidelity to parturition sites in sharks. *Molecular Ecology* **23**, 110-117.
- Fernando D (2012) A study of India's manta and mobula ray fishery. Manta Trust, Sri Lanka.
- Fernando D, Stevens G (2011) A study of Sri Lanka's manta and mobula ray fishery. Manta Trust, Sri Lanka.
- Feutry P, Kyne PM, Pillans RD, Chen X, Naylor G, Grewe PM (2014) Mitogenomics of the speartooth shark challenges ten years of Control Region sequencing. *BMC Evolutionary Biology* **14**, 232.

- Filippelli GM, Flores J-A (2009) From the warm Pliocene to the cold Pleistocene: A tale of two oceans. *Geology* **37**, 959-960.
- Fordyce RE (1980) Whale evolution and Oligocene Southern Ocean environments. *Palaeogeography, Palaeoclimatology, Palaeoecology* **31**, 319-336.
- Francis MP (2014) Survival and depth distribution of spinetail devilrays (*Mobula japanica*) released from purse-seine catches. In: NIWA Client Report No: WLG2014-2. Department of Conservation, New Zealand, Wellington, New Zealand.
- Funk WC, McKay JK, Hohenlohe PA, Allendorf FW (2012) Harnessing genomics for delineating Conservation Units. *Trends in Ecology and Evolution* **27**, 489-496.
- Furnestin J (1959) Hydrologie du Maroc Atlantique. *Revue des Travaux de l'Institut des Pêches Maritimes* **23**, 5-77.

G

- Gaida IH (1997) Population structure of the Pacific angel shark, *Squatina californica* (Squatini-formes: Squatinidae), around the California Channel Islands. *Copeia* **4**, 738-744.
- Galtier N, Nabholz B, Glemin S, Hurst GD (2009) Mitochondrial DNA as a marker of molecular diversity: a reappraisal. *Molecular Ecology* **18**, 4541-4550.
- Garcia VB, Lucifora LO, Myers RA (2008) The importance of habitat and life history to extinction risk in sharks, skates, rays and chimaeras. *Proceedings of the Royal Society B: Biological Sciences* **275**, 83-89.
- Gardner MG, Ward RD (1998) Population structure of the Australian gummy shark (*Mustelus antarcticus*) inferred from allozymes, mitochondrial DNA and vertebrae counts. *Marine and Freshwater Research* **49**, 733-745.
- Geraghty PT, Williamson JE, Macbeth WG, Wintner SP, Harry AV, Ovenden JR, Gillings MR (2013) Population expansion and genetic structure in *Carcharhinus brevipinna* in the southern Indo-Pacific. *PLoS One* **8**, e75169.
- Giles JL, Ovenden JR, AlMojil D, Garvilles E, Khametch K-o, Manjebraayakath H, Riginos C (2014) Extensive genetic population structure in the Indo-West Pacific spot-tail shark, *Carcharhinus sorrah*. *Bulletin of Marine Science* **90**, 427-454.
- Gill T (1908) The story of the devil fish. *Smithsonian Miscellaneous Collections* **52**, 155-180.
- Gillespie JH (2001) Is the population size of a species relevant to its evolution? *Evolution* **55**, 2161-2169.
- Gilman EL (2011) Bycatch governance and best practice mitigation technology in global tuna fisheries. *Marine Policy* **35**, 590-609.
- Glenn TC (2011) Field guide to Next-Generation DNA sequencers. *Molecular Ecology Resources* **11**, 759-769.
- Gomez-Gutierrez J, Tremblay N, Martinez-Gomez S, Robinson CJ, Del Angel-Rodriguez J, Rodriguez-Jaramillo C, Zavala-Hernandez C (2010) Biology of the subtropical sac-spawning euphausiid *Nyctiphanes simplex* in the northwestern seas of Mexico: vertical and horizontal distribution patterns and seasonal variability of brood size. *Deep-Sea Research Part II: Topical Studies in Oceanography* **57**, 606-615.
- Gonzalez-Isais M, Dominguez HM (2004) Comparative anatomy of the superfamily Myliobatoidea (Chondrichthyes) with some comments on phylogeny. *Journal of Morphol* **262**, 517-535.
- Gore MA, Rowat D, Hall J, Gell FR, Ormond RF (2008) Transatlantic migration and deep mid-ocean diving by basking shark. *Biology Letters* **4**, 395-398.
- Grada A, Weinbrecht K (2013) Next-Generation sequencing: methodology and application. *Journal of Investigative Dermatology* **133**, e11.
- Graham RT, Witt MJ, Castellanos DW, Remolina F, Maxwell S, Godley BJ, Hawkes LA (2012) Satel-

- lite tracking of manta rays highlights challenges to their conservation. PLoS One **7**, e36834.
- Graves JE (1998) Molecular insights into the population structures of cosmopolitan marine fishes. *Journal of Heredity* **89**, 427-437.
- Gubili C, Bilgin R, Kalkan E, Karhan SU, Jones CS, Sims DW, Kabasakal H, Martin AP, Noble LR (2011) Antipodean white sharks on a Mediterranean walkabout? Historical dispersal leads to genetic discontinuity and an endangered anomalous population. *Proceedings of the Royal Society B: Biological Sciences* **278**, 1679-1686.
- Gubili C, Sims DW, Verissimo A, Domenici P, Ellis J, Grigoriou P, Johnson AF, McHugh M, Neat F, Satta A, Scarcella G, Serra-Pereira B, Soldo A, Genner MJ, Griffiths AM (2014) A tale of two seas: contrasting patterns of population structure in the small-spotted catshark across Europe. *Royal Society Open Science* **1**, 140175-140175.
- H**
- Haddrill PR, Halligan DL, Tomaras D, Charlesworth B (2007) Reduced efficacy of selection in regions of the *Drosophila* genome that lack crossing over. *Genome Biology* **8**, R18.
- Hall M, Roman M (2013) Bycatches and non-tuna catches in the tropical tuna purse seine fisheries of the world.
- Han MV, Zmasek CM (2009) phyloXML: XML for evolutionary biology and comparative genomics. *BMC Bioinformatics* **10**, 356.
- Hasegawa T, Ando K, Mizundo K, Lukas R (2009) Coastal upwelling along the north coast of Papua New Guinea and SST Cooling over the Pacific Warm Pool: a case study for the 2002/03 El Niño event. *Journal of Oceanography* **65**, 817-833.
- Head ML, Kozak GM, Boughman JW (2013) Female mate preferences for male body size and shape promote sexual isolation in threespine sticklebacks. *Ecology and Evolution* **3**, 2183-2196.
- Heinrichs S, O'Malley MP, Medd H, Hilton P (2011) The global threat to *Manta* and *Mobula* rays.
- Henning W (1966) *Phylogenetic systematics* University of Illinois Press, Urbana, U.S.A.
- Herman J, Hovestadt-Euler M, Hovestadt DC, Stehmann M (2000) Contributions to the study of the comparative morphology of teeth and other relevant ichthyodorulites in living supra-specific taxa of Chondrichthyan fishes. Part B: Batomorphii 4c: Order: Rajiformes – Suborder Myliobatoidei – Superfamily Dasyatoidea – Family Dasyatidae – Subfamily Dasyatinae – Genus: *Urobatis*, Subfamily Potamotrygoninae – Genus: *Potamotrygon*, Superfamily Plesiobatoidea – Family Plesiobatidae – Genus: *Plesiobatis*, Superfamily Myliobatoidea – Family Myliobatidae – Subfamily Myliobatinae – Genera: *Aetobatus*, *Aetomylaeus*, *Myliobatis* and *Pteromylaeus*, Subfamily Rhinopterinae – Genus: *Rhinoptera* and Subfamily Mobulinae – Genera: *Manta* and *Mobula*. Addendum 1 to 4a: erratum to Genus *Pteroplatytrygon*. . *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique Entomologie* **70**, 5 - 67.
- Hershler R, Liu HP (2004) Taxonomic reappraisal of species assigned to the North American freshwater gastropod subgenus *Natricola* (Rissooidea: Hydrobiidae). *Veliger* **47**, 66-81.
- Hey J (2005) On the number of New World founders: a population genetic portrait of the peopling of the Americas. *PLoS Biology* **3**, e193.
- Hey J, Nielsen R (2004) Multilocus methods for estimating population sizes, migration rates and divergence time, with applications to the divergence of *Drosophila pseudoobscura* and *D. persimilis*. *Genetics* **167**, 747-760.
- Hey J, Nielsen R (2007) Integration within the Felsenstein equation for improved Markov chain Monte Carlo methods in population genetics. *Proceedings of the National Academy of Sciences* **104**, 2785-2790.
- Hillis DM, Pollock DD, McGuire JA, Zwickl DJ (2003) Is sparse taxon sampling a problem for phylogenetic inference? *Systematic Biology* **52**, 124-126.

- Hinojosa-Alvarez S, Díaz-Jaimes P, Marcet-Houben M, Gabaldón T (2014) The complete mitochondrial genome of the giant manta ray, *Manta birostris*. *Mitochondrial DNA* **0**, 1-2.
- Hoarau G, Coyer JA, Stam WT, Olsen JL (2006) A fast and inexpensive DNA extraction/purification protocol for brown macroalgae. *Molecular Ecology Notes* **7**, 191-193.
- Hoarau G, Holla S, Lescasse R, Stam WT, Olsen JL (2002) Heteroplasmy and evidence for recombination in the mitochondrial Control Region of the flatfish *Platichthys flesus*. *Molecular Biology and Evolution* **19**, 2261-2264.
- Hoelzel AR, Natoli A, Dahlheim ME, Olavarria C, Baird RW, Black NA (2002) Low worldwide genetic diversity in the killer whale (*Orcinus orca*): implications for demographic history. *Proceedings of the Royal Society B: Biological Sciences* **269**, 1467-1473.
- Hoelzel AR, Shivji MS, Magnussen J, Francis MP (2006) Low worldwide genetic diversity in the basking shark (*Cetorhinus maximus*). *Biology Letters* **2**, 639-642.
- Hoening JM, Gruber SH (1990) Life-history patterns in the Elasmobranchs: implications for fisheries management NOAA Technical report
- Holcer D, Lazar B, Mackelworth P, Fortuna CM (2013) Rare or just unknown? The occurrence of the giant devil ray (*Mobula mobular*) in the Adriatic Sea. *Journal of Applied Ichthyology* **29**, 139-144.
- Homma K, Maruyama T, Itoh T, Ishihara H, Uchida S (1999) Biology of the manta ray, *Manta birostris* Walbaum, in the Indo-Pacific. In: Indo-Pacific fish biology: proceedings of the fifth international conference on Indo-Pacific fishes, Noumea, 1997. eds. Seret B, Sire J), pp. 209-216. Ichthyological Society of France, Paris, France.
- Huelsenbeck J, Ronquist F (2001) MRBAYES: bayesian inference of phylogenetic trees. *Bioinformatics* **17**, 754-755.
- Hueter RE, Heupel MR, Heist EJ, Keeney DB (2004) Evidence of philopatry in sharks and implications for the management of shark fisheries. *Journal of Northwest Atlantic Fishery Science* **37**, 239-247.
- Hunter E, Buckley AA, Stewart C, Metcalfe JD (2005) Migratory behaviour of the thornback ray, *Raja clavata*, in the southern North Sea. *Journal of the Marine Biological Association of the UK* **85**.
- Hutchings J, Myers R (2012) Life-history correlates of extinction risk and recovery potential. *Ecological Applications* **22**, 1061-1067.
- I**
- IATTC (2012) Tunas and billfishes in the Eastern Pacific Ocean in 2011. Fishery Status Report No. **10**.
- Ikema T, Bryant RG, Bigg GR (2013) Environmental controls at multiple scales for the western Pacific: an Okinawan case study. *Estuarine, Coastal and Shelf Science* **128**, 52-63.
- Irwin DE (2002) Phylogeographic breaks without geographic barriers to gene flow. *Evolution* **56**, 2383-2394.
- Ishikawa S, Oda M (2007) Reconstruction of Indian monsoon variability over the past 230,000 years: Planktic foraminiferal evidence from the NW Arabian Sea open-ocean upwelling area. *Marine Micropaleontology* **63**, 143-154.
- IUCN (2012) The IUCN Red List of Threatened Species Version. 2011.2.
- J**
- Jackson AM, Ambariyanto, Erdmann MV, Toha AHA, Stevens LA, Barber PH (2014) Phylogeography of commercial tuna and mackerel in the Indonesian Archipelago. *Bulletin of Marine Science* **90**, 471-492.
- Jarne P, Lagoda PJL (1996) Microsatellites, from molecules to populations and back. *Trends in*

- Ecology & Evolution **11**, 424-429.
- Jones E, Francis MP (2012) Protected rays – occurrence and development of mitigation methods in the New Zealand tuna purse seine fishery. Client Report No. WEG2012-49. Department of Conservation, New Zealand, Auckland, New Zealand.
- Jorgensen SJ, Reeb CA, Chapple TK, Anderson S, Perle C, Van Sommeran SR, Fritz-Cope C, Brown AC, Klimley AP, Block BA (2010) Philopatry and migration of Pacific white sharks. *Proceedings of the Royal Society B: Biological Sciences* **277**, 679-688.
- Jost L (2008) Gst and its relatives do not measure differentiation. *Molecular Ecology* **17**, 4015-4026.
- K**
- Karl SA, Castro ALF, Garla RC (2011) Population genetics of the nurse shark (*Ginglymostoma cirratum*) in the western Atlantic. *Marine Biology* **159**, 489-498.
- Karl SA, Castro ALF, Lopez JA, Charvet P, Burgess GH (2010) Phylogeography and conservation of the bull shark (*Carcharhinus leucas*) inferred from mitochondrial and microsatellite DNA. *Conservation Genetics* **12**, 371-382.
- Kashiwagi T, Marshall AD, Bennett MB, Ovenden JR (2012) The genetic signature of recent speciation in manta rays (*Manta alfredi* and *M. birostris*). *Molecular Phylogenetics and Evolution* **64**, 212-218.
- Keeney DB, Heist EJ (2006) Worldwide phylogeography of the blacktip shark (*Carcharhinus limbatus*) inferred from mitochondrial DNA reveals isolation of western Atlantic populations coupled with recent Pacific dispersal. *Molecular Ecology* **15**, 3669-3679.
- Keeney DB, Heupel M, Hueter RE, Heist EJ (2003) Genetic heterogeneity among blacktip shark, *Carcharhinus limbatus*, continental nurseries along the U.S. Atlantic and Gulf of Mexico. *Marine Biology* **143**, 1039-1046.
- Keeney DB, Heupel MR, Hueter RE, Heist EJ (2005) Microsatellite and mitochondrial DNA analyses of the genetic structure of blacktip shark (*Carcharhinus limbatus*) nurseries in the northwestern Atlantic, Gulf of Mexico, and Caribbean Sea. *Molecular Ecology* **14**, 1911-1923.
- Kelchner SA, Thomas MA (2007) Model use in phylogenetics: nine key questions. *Trends in Ecology and Evolution* **22**, 87-94.
- Keller LF, Waller DM (2002) Inbreeding effects in wild populations. *Trends in Ecology and Evolution* **17**, 230-241.
- Kollias S, Poortvliet M, Smolina I, Hoarau G (2015) A quick and easy method of mitochondrial genome capture for NGS applications. *Conservation Genetics Resources* **7**, 345-348.
- Koutavas A, Lynch-Stieglitz J, Marchitto TM, Sachs JP (2002) El Niño-like pattern in ice age tropical Pacific sea surface temperature. *Science* **297**, 226-230.
- Koutavas A, Lynch-Stieglitz J (2003) Glacial-interglacial dynamics of the eastern equatorial Pacific cold tongue-Intertropical Convergence Zone system reconstructed from oxygen isotope records. *Paleoceanography* **18**.
- Krefft G, Stehmann M (1973) *Mobulidae*. In: Check-list of the fishes of the North-eastern Atlantic and of the Mediterranean (CLOFNAM) (eds. Hureau JC, Monod T), p. 77. Unesco, Paris.
- Krishna KM (2011) Satellite surveillance of upwelling along the east coast of India. *Marine Geodesy* **34**, 181-187.
- Kubo T, Iwasa Y (1995) Inferring the rates of branching and extinction from molecular phylogenies. *Evolution* **49**, 694-704.
- L**
- LaFond EC (1954) On upwelling and sinking off the east coast of India. *Andhra University Mem-*

- oires in Oceanography **1**, 117-121.
- LaFond EC (1957) Oceanographic studies in the Bay of Bengal. Proceedings of the Indian Academy of Science, Section B **46**, 1-46.
- Lanfear R, Calcott B, Ho SY, Guindon S (2012) Partitionfinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. Molecular Biology and Evolution **29**, 1695-1701.
- Last PR, Stevens JD (1994) Sharks and Rays of Australia CSIRO Publishing, Australia.
- Laurito Mora CA (1999) Los selaceos fosiles de la localidad de Alto Guayacan (y otros ictiolitos asociados): Mioceno Superior-Plioceno Inferior de Limón Guila Imprenta, San José, Costa Rica.
- Le Port A, Lavery S (2012) Population structure and phylogeography of the short-tailed stingray, *Dasyatis brevicaudata* (Hutton 1875), in the Southern Hemisphere. Journal of Heredity **103**, 174-185.
- Lessios HA (2008) The great American schism: divergence of marine organisms after the rise of the Central American Isthmus. Annual Review of Ecology, Evolution, and Systematics **39**, 63-91.
- Lessios HA, Kessing BD, Pearse JS (2001) Population structure and speciation in tropical seas: global phylogeography of the sea urchin *Diadema*. Evolution **55**, 955-975.
- Lewallen EA, Anderson TW, Bohonak AJ (2007) Genetic structure of leopard shark (*Triakis semifasciata*) populations in California waters. Marine Biology **152**, 599-609.
- Lewison R, Crowder L, Read A, Freeman S (2004) Understanding impacts of fisheries bycatch on marine megafauna. Trends in Ecology and Evolution **19**, 598-604.
- Li N, Song N, Cheng G-P, Gao T-X (2013) Genetic diversity and population structure of the red stingray, *Dasyatis akajei* inferred by AFLP marker. Biochemical Systematics and Ecology **51**, 130-137.
- Li YC, Korol AB, Fahima T, Beiles A, Nevo E (2002) Microsatellites: genomic distribution, putative functions and mutational mechanisms: a review. Molecular Ecology **11**, 2453-2465.
- Librado P, Rozas R (2009) DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. Bioinformatics **25**, 1451-1452.
- Lloyd RE (1908) On two species of eagle-rays with notes on the skull of the genus *Ceratoptera*. Records Indian Museum (Calcutta) **2**, 175-180.
- Lovejoy NR (1996) Systematics of myliobatoid elasmobranchs: with emphasis on the phylogeny and historical biogeography of neotropical freshwater stingrays (Potamotrygonidae: Rajiformes). Zoological Journal of the Linnean Society **117**, 207-257.
- Luiz OJ, Balboni AP, Kodja G, Andrade M, Marum H (2009) Seasonal occurrences of *Manta birostris* (Chondrichthyes: Mobulidae) in southeastern Brazil. Ichthyological Research **56**, 96-99.
- Lunt DH, Whipple LE, Hyman BC (1998) Mitochondrial DNA variable number tandem repeats (VNTRs): utility and problems in molecular ecology. Molecular Ecology **7**, 1441-1455.
- Lutjeharms JRE, Machu E (2000) An upwelling cell inshore of the East Madagascar Current. Deep Sea Research Part I: Oceanographic Research Papers **47**, 2405-2411.

M

- Ma W, Chai F, Xiu P, Xue H, Tian J (2013) Modeling the long-term variability of phytoplankton functional groups and primary productivity in the South China Sea. Journal of Oceanography **69**, 527-544.
- Marlow JR, Lange CB, Wefer G, Rosell-Mele A (2000) Upwelling intensification as part of the Pliocene-Pleistocene climate transition. Science **290**, 2288-2291.
- Marshall AD, Bennett MB (2010) Reproductive ecology of the reef manta ray *Manta alfredi* in southern Mozambique. Journal of Fish Biology **77**, 169-190.

- Marshall AD, Compagno LJ, Leonard JV, Bennett MB (2009) Redescription of the genus *Manta* with resurrection of *Manta alfredi* (Krefft, 1868) (Chondrichthyes; Myliobatoidei; Mobulidae). *Zootaxa*, 1-28.
- Marshall AD, Kashiwagi T, Bennett MB, Deakos MH, Stevens G, McGregor F, Clark TB, Ishihara H, Sato K (2011) *Manta alfredi*. In: The IUCN Red List of Threatened Species. Version 2013.2. www.iucnredlist.org. Accessed on 22 May 2014.
- Martin AP (1995) Mitochondrial DNA sequence evolution in sharks: rates, patterns and phylogenetic inference. *Molecular Biology and Evolution* **12**, 1114-1123.
- Martin AP (1999) Substitution rates of organelle and nuclear genes in sharks: implicating metabolic rate (again). *Molecular Biology and Evolution* **16**, 996-1002.
- Martin AP, Naylor GJ, Palumbi SR (1992) Rates of mitochondrial DNA evolution in sharks are slow compared with mammals. *Nature* **357**.
- Martin AP, Palumbi SR (1993) Body size, metabolic rate, generation time, and the molecular clock. *Proceedings of the National Academy of Sciences U.S.A* **90**.
- Martinez P, Gonzalez EG, Castilho R, Zardoya R (2006) Genetic diversity and historical demography of Atlantic bigeye tuna (*Thunnus obesus*). *Molecular Phylogenetics and Evolution* **39**, 404-416.
- Mason VC, Li G, Helgen KM, Murphy WJ (2011) Efficient cross-species capture hybridization and Next-Generation sequencing of mitochondrial genomes from noninvasively sampled museum specimens. *Genome Research* **21**, 1695-1704.
- Mayr E (1963) *Animal species and evolution* Belknap Press of Harvard University Press, Cambridge, USA.
- Mazeika PA (1967) Thermal domes in the eastern tropical Atlantic. *Limnology and Oceanography* **12**, 537-539.
- McCauley DJ, DeSalles PA, Young HS, Papastamatiou YP, Caselle JE, Deakos MH, Gardner JPA, Garton DW, Collen JD, Micheli F (2014) Reliance of mobile species on sensitive habitats: a case study of manta rays (*Manta alfredi*) and lagoons. *Marine Biology* DOI: 10.1007/s00227-014-2478-7.
- McEachran JD, Aschliman NC (2004) Phylogeny of Batoidea. In: *Biology of sharks and their relatives* (eds. Carrier JC, Musick JA, Heithaus MR). CRC Press, Boca Raton, FL.
- McEachran JD, Dunn KA, Miyake T (1996) Interrelationships of the batoid fishes (Chondrichthyes, Batoidea). In: *Interrelationships of Fishes* (eds. Stiassny MLJ, Parenti LR, Johnson GD), pp. 63-82. Academic Press., San Diego, CA.
- Meirmans PG, Hedrick PW (2011) Assessing population structure: F_{ST} and related measures. *Molecular Ecology Resources* **11**, 5-18.
- Meirmans PG, Van Tienderen PH (2004) Genotype and Genodive: two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes* **4**, 792-794.
- Mendonça FF, Oliveira C, Gadig OBF, Foresti F (2011) Phylogeography and genetic population structure of Caribbean sharpnose shark *Rhizoprionodon porosus*. *Reviews in Fish Biology and Fisheries* **21**, 799-814.
- Mendonça FF, Oliveira C, Gadig OBF, Foresti F (2013) Diversity and genetic population structure of the Brazilian sharpnose shark *Rhizoprionodon lalandii*. *Aquatic Conservation: Marine and Freshwater Ecosystems* **23**, 850-857.
- Millesi AC, Oddone MC (2003) Primer registro de *Manta birostris* (Donndurff 1798) (Batoidea: Mobulidae) en el Rio de la Plata, Uruguay. *Gayana* **67**, 127-113-.
- Miya M, Takeshima H, Endo H, Ishiguro NB, Inoue JG, Mukai T, Satoh TP, Yamaguchi M, Kawaguchi A, Mabuchi K, Shirai SM, Nishida M (2003) Major patterns of higher teleostean phylogenies: a new perspective based on 100 complete mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* **26**, 121-138.

- Mohanraj G, Rajapackiam S, Mohan S, Batcha H, Gomathy S (2009) Status of elasmobranchs fishery in Chennai, India. *Asian Fisheries Science* **22**, 607-615.
- Molony B (2005) Estimates of the mortality of non-target species with an initial focus on seabirds, turtles and sharks. In: 1st Meeting of the Scientific Committee of the Western and Central Pacific Fisheries Commission. Report No. WCPFC-SC1 EB WP-1. Western and Central Pacific Fisheries Commission (WCPFC), Noumea, New Caledonia.
- Molony B (2008) Fisheries biology and ecology of highly migratory species that commonly interact with industrialised longline and purse-seine fisheries in the western and central Pacific. In: Scientific Committee Fourth Regular Session, Report No. WCPFC-SC4-2008/EB-IP-6. Western and Central Pacific Fisheries Commission (WCPFC), Port Moresby, Papua New Guinea.
- Moore WS (1995) Inferring phylogenies from mtDNA variation: mitochondrial-gene trees versus nuclear-gene trees. *Evolution* **49**, 718-726.
- Morin PA, Archer FI, Foote AD, Vilstrup J, Allen EE, Wade P, Durban J, Parsons K, Pitman R, Li L, Bouffard P, Abel Nielsen SC, Rasmussen M, Willerslev E, Gilbert MT, Harkins T (2010) Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Research* **20**, 908-916.
- Moritz C (1994a) Applications of mitochondrial DNA analysis in conservation: a critical review. *Molecular Ecology* **3**, 401-411.
- Moritz C (1994b) Defining 'Evolutionary Significant Units' for conservation. *TREE* **9**, 373-375.
- Morlon H, Parsons TL, Plotkin JB (2011) Reconciling molecular phylogenies with the fossil record. *Proceedings of the National Academy of Sciences U.S.A* **108**, 16327-16332.
- Morrison WR, Lohr JL, Duchon P, Wilches R, Trujillo D, Mair M, Renner SS (2009) The impact of taxonomic change on conservation: does it kill, can it save, or is it just irrelevant? *Biological Conservation* **142**, 3201-3206.
- Mourier J, Planes S (2013) Direct genetic evidence for reproductive philopatry and associated fine-scale migrations in female blacktip reef sharks (*Carcharhinus melanopterus*) in French Polynesia. *Mol Ecol* **22**, 201-214.
- Müller J, Henle J (1841) *Systematische beschreibung der plagiostomen* Verlag von Veit & Co., Berlin.
- Müller K (2005) SeqState - primer design and sequence statistics for phylogenetic DNA data sets. *Applied Bioinformatics* **4**, 65-69.
- Myers RA, Worm B (2005) Extinction, survival or recovery of large predatory fishes. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **360**, 13-20.

N

- Nabholz B, Glemin S, Galtier N (2008) Strong variations of mitochondrial mutation rate across mammals - the longevity hypothesis. *Molecular Biology and Evolution* **25**, 120-130.
- Nance HA, Klimley P, Galván-Magaña F, Martínez-Ortiz J, Marko PB (2011) Demographic processes underlying subtle patterns of population structure in the scalloped hammerhead shark, *Sphyrna lewini*. *PLoS One* **6**, e21459.
- Naylor GJP, Caira JN, Jensen K, Rosana KAM, Straube N, Lakner C (2012) Elasmobranch phylogeny: a mitochondrial estimate based on 595 species. In: *Biology of sharks and their relatives* (ed. Carrier JCMJAHMR). CRC Press, Boca Raton.
- Nee S, May RM, Harvey PH (1994) The reconstructed evolutionary process. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **344**, 305-311.
- Nishida K (1990) *Phylogeny of the suborder Myliobatidoidei*, Hokkaido University.
- Notarbartolo di Scara G, Serena F, Mancusi C (2007) Giant devil ray *Mobula mobular* (Bonnaterre, 1788). In: *Overview of the conservation status of cartilaginous fishes (Chondrich-*

- thyans) in the Mediterranean Sea (eds. Cavanagh RD, Gibson C), p. 42. The World Conservation Union (IUCN), Gland, Switzerland.
- Notarbartolo di Sciara G (1987) A revisionary study of the genus *Mobula* Rafinesque, 1810 (Chondrichthyes: Mobulidae) with the description of a new species. *Zoological Journal of the Linnean Society* **91**, 1-91.
- Notarbartolo di Sciara G (1988) Natural history of the rays of the genus *Mobula* in the Gulf of California. *Fishery Bulletin* **86**.
- Notarbartolo di Sciara G, Serena F, Mancusi C (2011) *Mobula mobular*. In: IUCN Red List of Threatened Species.
- Nylander JA, Wilgenbusch JC, Warren DL, Swofford DL (2008) AWTY (are we there yet?): a system for graphical exploration of MCMC convergence in Bayesian phylogenetics. *Bioinformatics* **24**, 581-583.
- O**
- O'Connell M, Wright JM (1997) Microsatellite DNA in fishes. *Reviews in Fish Biology and Fisheries* **7**, 331-363.
- O'Malley MP, Lee-Brooks K, Medd HB (2013) The global economic impact of manta ray watching tourism. *PLoS One* **8**, e65051.
- O'Brien SM, Gallucci VF, Hauser L (2012) Effects of species biology on the historical demography of sharks and their implications for likely consequences of contemporary climate change. *Conservation Genetics* **14**, 125-144.
- Olson PD, Caira JN, Jensen K, Overstreet RM, Palm HW, Beveridge I (2010) Evolution of the trypanorhynch tapeworms: parasite phylogeny supports independent lineages of sharks and rays. *International Journal of Parasitology* **40**, 223-242.
- Ortiz M, Prince ED, Serafy JE, Holts DB, Davy KB, Pepperell JG, Lowry MB, Holdsworth JC (2003) Global overview of the major constituent-based billfish tagging programs and their results since 1954. *Marine and Freshwater Research* **54**, 489-507.
- Ovenden JR, Kashiwagi T, Broderick D, Giles J, Salini J (2009) The extent of population genetic subdivision differs among four co-distributed shark species in the Indo-Australian archipelago. *BMC Evolutionary Biology* **9**, 40.
- Ovenden JR, Morgan JAT, Street R, Tobin A, Simpfendorfer C, Macbeth W, Welch D (2011) Negligible evidence for regional genetic population structure for two shark species *Rhizoprionodon acutus* (Rüppell, 1837) and *Sphyrna lewini* (Griffith & Smith, 1834) with contrasting biology. *Marine Biology* **158**, 1497-1509.
- Owens IPF, Bennett PM (2000) Ecological basis of extinction risk in birds: habitat loss versus human persecution and introduced predators. *Proceedings of the National Academy of Sciences* **97**, 12144.
- P**
- Paetkau D, Slade R, Burden M, Estoup A (2004) Genetic assignment methods for the direct, real-time estimation of migration rate: a simulation-based exploration of accuracy and power. *Molecular Ecology* **13**, 55-65.
- Paig-Tran EW, Kleinteich T, Summers AP (2013) The filter pads and filtration mechanisms of the devil rays: variation at macro and microscopic scales. *Journal of Morphology* **274**, 1026-1043.
- Palsboll PJ, Berube M, Allendorf FW (2007) Identification of management units using population genetic data. *Trends in Ecology and Evolution* **22**, 11-16.
- Palumbi SR (1994) Genetic divergence, reproductive isolation, and marine speciation. *Annual Review in Ecology and Systematics* **25**, 547-572.

- Pannell JR (2003) Coalescence in a metapopulation with recurrent local extinction and recolonization. *Evolution* **57**, 949-961.
- Paradis E, Claude J, Strimmer K (2004) APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* **20**, 289-290.
- Pardini AT, Jones CS, Noble LR, Kreiser B, Malcolm H, Bruce BD, D SJ, Cliff G, Scholl MC, Francis M, Duffy CAJ, Martin AP (2001) Sex-biased dispersal of great white sharks. *Nature* **412**, 139-140.
- Pastene LA, Goto M, Kanda N, Zerbini AN, Kerem D, Watanabe K, Bessho Y, Hasegawa M, Nielsen R, Larsen F, Palsboll PJ (2007) Radiation and speciation of pelagic organisms during periods of global warming: the case of the common minke whale, *Balaenoptera acutorostrata*. *Molecular Ecology* **16**, 1481-1495.
- Patrick A, Thunell RC (1997) Tropical Pacific sea surface temperatures and upper water column thermal structure during the Last Glacial Maximum. *Paleoceanography* **12**, 649-657.
- Paulin CD, Habib G, Carey CL, Swanson PM, Voss GJ (1982) New records of *Mobula japonica* and *Masturus lanceolatus*, and further records of *Luvaris imperialis*. *New Zealand Journal of Marine and Freshwater Research* **16**, 11-17.
- Pedersen TF (1983) Increased productivity in the eastern equatorial Pacific during the Last Glacial Maximum (19,000 to 14,000 yr B.P.). *Geology* **11**, 16-19.
- Peeters FJC, Acheson R, Brummer GJA, de Ruijter WPM, Schneider RR, Ganssen GM, Ufkes E, Kroon D (2004) Vigorous exchange between the Indian and Atlantic oceans at the end of the past five glacial periods. *Nature* **430**, 661-665.
- Pellegrin J (1901) Sur une raie cornue gigantesque pechee à Oran. *Bulletin du Museum National d'Histoire Naturelle* **7**, 327-328.
- Pennings P, Achenbach A, Foitzik S (2011) Data from: Similar evolutionary potentials in an obligate ant parasite and its two host species, Dryad Digital Repository.
- Pereyra S, García G, Miller P, Oviedo S, Domingo A (2010) Low genetic diversity and population structure of the narrownose shark (*Mustelus schmitti*). *Fisheries Research* **106**, 468-473.
- Pfeil FH (1981) Eine nektonische Fischfauna aus dem unteroligozänen Schöneck-Fischschiefer des Galon-Grabens in Oberbayern. *Geologica Bavarica* **82**, 357-388.
- Philippi RA (1892) La conductiofauna de las regiones de Chile y California-Oregon: comparacion preliminar, Viña del Mar, Chile.
- Pianet R, Nordstrom V, Damiano A, Delgado de Molina A, Ariz J, Sabate I, Kouassi Y, Sow F (2010) Statistiques de la pêche thonière Européenne et assimilée dans l'Océan Atlantique durant la période 1991-2008. *Collective Volume Scientific Papers ICCAT* **65**, 561-591.
- Picot L, Becker D, Cavin L, Pirkenseer C, Lapaire F, Rauber G, Hochuli PA, J-P. B (2008) Sédimentologie et paléontologie des paléoenvironnements côtiers Rupéliens de la Molasse marine rhénane dans le Jura suisse. *Swiss Journal of Geosciences* **101**, 483-513.
- Piela C, Lyle M, Marcantonio F, Baldauf J, Olivarez Lyle A (2012) Biogenic sedimentation in the equatorial Pacific: carbon cycling and paleoproduction, 12-24 Ma. *Paleoceanography* **27**.
- Pillans B, Chappell J, Naish TR (1998) A review of the Milankovitch climatic beat: template for Plio-Pleistocene sea-level changes and sequence stratigraphy. *Sedimentary Geology* **122**, 5-21.
- Piry S, Alapetite A, Cornuet J-M, Paetkau D, Baudouin L, Estoup A (2004) GENECLASS2: a software for genetic assignment and first-generation migrant detection. *Journal of Heredity* **95**, 536-539.
- Planes S, Doherty PJ, Bernardi G (2001) Strong genetic divergence among populations of a marine fish with limited dispersal, *Acanthochromis polyacanthus*, within the Great Barrier Reef and the Coral Sea. *Evolution* **55**, 2263-2273.
- Plank SM, Lowe CG, Feldheim KA, Wilson RR, Jr., Brusslan JA (2010) Population genetic structure

- of the round stingray *Urobatis halleri* (Elasmobranchii: Rajiformes) in southern California and the Gulf of California. *Journal of Fish Biology* **22**, 329-340.
- Poisson F, Séret B, Vernet A-L, Goujon M, Dagorn L (2014) Collaborative research: development of a manual on elasmobranch handling and release best practices in tropical tuna purse-seine fisheries. *Marine Policy* **44**, 312-320.
- Pollock DD, Zwickl DJ, McGuire JA, Hillis DM (2002) Increased taxon sampling is advantageous for phylogenetic inference. *Systematic Biology* **51**, 664-671.
- Polovina JJ, Dunne JP, Woodworth PA, Howell EA (2011) Projected expansion of the subtropical biome and contraction of the temperate and equatorial upwelling biomes in the North Pacific under global warming. *ICES Journal of Marine Science* **68**, 986-995.
- Poortvliet M, Galván-Magaña F, Bernardi G, Croll DA, Olsen JL (2011) Isolation and characterization of twelve microsatellite loci for the Japanese Devilray (*Mobula japanica*). *Conservation Genetics Resources* **3**, 733-735.
- Poortvliet M, Hoarau G (2013) The complete mitochondrial genome of the spinetail devilray, *Mobula japanica*. *Mitochondrial DNA* **24**, 28-30.
- Poortvliet M, Hoarau G, Croll DA, Bernardi G, Newton KM, O'Sullivan J, Fernando D, Stevens G, Magana FG, Seret B, Olsen JL (in preparation) Influences of past climatic change on historical population structure and demography of the spinetail devilray, *Mobula japanica*, based on mitogenomes and nuclear DNA.
- Poortvliet M, Olsen JL, Croll DA, Bernardi G, Newton KM, Kollias S, O'Sullivan J, Fernando D, Stevens G, Galván-Magaña F, Seret B, Wintner S, Hoarau G (2015) A dated molecular phylogeny of manta and devil rays (Mobulidae) based on mitogenome and nuclear sequences. *Molecular Phylogenetics and Evolution* **83**, 72-85.
- Portnoy DS, Hollenbeck CM, Belcher CN, Driggers WB, 3rd, Frazier BS, Gelsleichter J, Grubbs RD, Gold JR (2014) Contemporary population structure and post-glacial genetic demography in a migratory marine species, the blacknose shark, *Carcharhinus acronotus*. *Molecular Ecology* **23**, 5480-5495.
- Portnoy DS, McDowell JR, Heist EJ, Musick JA, Graves JE (2010) World phylogeography and male-mediated gene flow in the sandbar shark, *Carcharhinus plumbeus*. *Molecular Ecology* **19**, 1994-2010.
- Posada D (2008) jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* **25**, 1253-1256.
- Presgraves DC (2005) Recombination enhances protein adaptation in *Drosophila melanogaster*. *Current Biology* **15**, 1651-1656.
- Purdy RW, Schneider VP, Applegate SP, McLellan JH, Meyer RL, Slaughter BH (2001) The Neogene sharks, rays, and bony fishes from Lee Creek Mine, Aurora, North Carolina. In: *Geology and paleontology of the Lee Creek Mine, North Carolina, III* (eds. Clayton ER, Bohaska DJ), pp. 71-202. Smithsonian Institution Press, Washington.

R

- Rafinesque-Schmaltz CS (1810) *Indice d'ittologia siciliana ossia catalogo metodico dei nomi latini, italiani, e siciliani dei pesci, che si rinvencono in Sicilia, Messina Presso Giovanni del Nobolo, Messina, Italy.*
- Rajapackiam S, Mohan S, Rudramurthy N (2007) Utilization of gill rakers of lesser devil ray *Mobula diabolus* - a new fish byproduct. *Marine Fisheries Information Service, Technical and Extension Series* **191**, 22-23.
- Raje SG, Zacharia PU, Gopal-Raje S (2009) Investigations on fishery and biology of nine species of rays in Mumbai waters. *Indian Journal of Fisheries* **56**, 95-101.
- Rama Sastry AA, Myrland P (1959) Distribution of temperature, salinity and density in the

- Arabian Sea along the south Malabar coast (South India) during the post-monsoon season. *Indian Journal of Fisheries* **6**, 223-255.
- Rambaut A, Drummond AJ (2009) Tracer v1.5. <http://tree.bio.ed.ac.uk/software/tracer/>
- Rand DM (1993) Endotherms, Ectotherms, and Mitochondrial Genome-Size Variation. *Journal of Molecular Evolution* **37**, 281-295.
- Rannala B, Mountain J (1997) Detecting immigration by using multilocus genotypes. *Proceedings of the National Academy of Sciences* **94**, 9197-9201.
- Ravelo AC, Andreasen DH, Lyle M, Lyle AO, Wara MW (2004) Regional climate shifts caused by gradual global cooling in the Pliocene epoch. *Nature* **429**.
- Reich D, Green RE, Kircher M, Krause J, Patterson N, Durand EY, Viola B, Briggs AW, Stenzel U, Johnson PL, Maricic T, Good JM, Marques-Bonet T, Alkan C, Fu Q, Mallick S, Li H, Meyer M, Eichler EE, Stoneking M, Richards M, Talamo S, Shunkov MV, Dereviako AP, Hublin JJ, Kelso J, Slatkin M, Paabo S (2010) Genetic history of an archaic hominin group from Denisova Cave in Siberia. *Nature* **468**, 1053-1060.
- Revell LJ, Harmon LJ, Glor RE (2005) Underparameterized model of sequence evolution leads to bias in the estimation of diversification rates from molecular phylogenies. *Systematic Biology* **54**, 973-983.
- Riccioni G, Landi M, Ferrara G, Milano I, Cariani A, Zane L, Sella M, Barbujani G, Tinti F (2010) Spatio-temporal population structuring and genetic diversity retention in depleted Atlantic bluefin tuna of the Mediterranean Sea. *Proceedings of the National Academy of Science U.S.A* **107**, 2102-2107.
- Richards VP, Henning M, Witzell W, Shivji MS (2009) Species delineation and evolutionary history of the globally distributed spotted eagle ray (*Aetobatus narinari*). *Journal of Heredity* **100**, 273-283.
- Rivers MC, Brummitt NA, Nic Lughadha E, Meagher TR (2014) Do species conservation assessments capture genetic diversity? *Global Ecology and Conservation* **2**, 81-87.
- Rohfritsch A, Borsa P (2005) Genetic structure of Indian scad mackerel *Decapterus russelli*: Pleistocene vicariance and secondary contact in the Central Indo-West Pacific Seas. *Heredity* **95**, 315-326.
- Rohner C, Pierce S, Marshall A, Weeks S, Bennett M, Richardson A (2013) Trends in sightings and environmental influences on a coastal aggregation of manta rays and whale sharks. *Marine Ecology Progress Series* **482**, 153-168.
- Romanov EV (2002) Bycatch in the tuna purse-seine fisheries of the western Indian Ocean. *Fishery Bulletin* **100**, 90-105.
- Rooker JR, Secor DH, de Metrio G, Schloesser R, Block BA, Neilson JD (2008) Natal homing and connectivity in Atlantic Bluefin tuna populations. *Science* **322**, 742-744.
- Roosevelt T (1917) Harpooning devilfish. *Scribner's Magazine* **62**, 293-305.
- Rosel PE, Block BA (1996) Mitochondrial Control Region variability and global population structure in the swordfish, *Xiphias gladius*. *Marine Biology* **125**, 11-22.
- Ross MG, Russ C, Costello M, Hollinger A, Lennon NJ, Hegarty R, Nusbaum C, Jaffe DB (2013) Characterizing and measuring bias in sequence data. *Genome Biology* **14**, R51.
- Roughan M, Middleton JH (2002) A comparison of observed upwelling mechanisms off the east coast of Australia. *Continental Shelf Research* **22**, 2551-2572.
- Ryman N, Leimar O (2009) Gst is still a useful measure of genetic differentiation - a comment on Jost's D. *Molecular Ecology* **18**, 2084-2087.

S

- Sainudiin R, Durrett RT, Aquadro CF, Nielsen R (2004) Microsatellite mutation models: insights from a comparison of humans and chimpanzees. *Genetics* **168**, 383-395.

- Sandoval-Castillo J, Rocha-Olivares A (2011) Deep mitochondrial divergence in Baja California populations of an aquilopelagic elasmobranch: the golden cownose ray. *Journal of Heredity* **102**, 269-274.
- Sandoval-Castillo J, Rocha-Olivares A, Villavicencio-Garayzar C, Balart E (2004) Cryptic isolation of Gulf of California shovelnose guitarfish evidenced by mitochondrial DNA. *Marine Biology* **145**, 983-988.
- Schaefer KM, Fuller DW, Block BA (2007) Movements, behavior, and habitat utilization of yellowfin tuna (*Thunnus albacares*) in the northeastern Pacific Ocean, ascertained through archival tag data. *Marine Biology* **152**, 503-525.
- Schluessel V, Broderick D, Collin SP, Ovenden JR (2010) Evidence for extensive population structure in the white-spotted eagle ray within the Indo-Pacific inferred from mitochondrial gene sequences. *Journal of Zoology* **281**, 46-55.
- Schmittner A (2005) Decline of the marine ecosystem caused by a reduction in the Atlantic overturning circulations. *Nature* **434**, 628-633.
- Schrey AW, Heist EJ (2003) Microsatellite analysis of population structure in the shortfin mako (*Isurus oxyrinchus*). *Canadian Journal of Fisheries and Aquatic Sciences* **60**, 670-675.
- Schuelke M (2000) An economic method for the fluorescent labeling of PCR fragments. *Nature* **18**, 233-234.
- Schultz JK, Feldheim KA, Gruber SH, Ashley MV, McGovern TM, Bowen BW (2008) Global phylogeography and seascape genetics of the lemon sharks (genus *Negaprion*). *Molecular Ecology* **17**, 5336-5348.
- Serrano-López JN (2009) Estudio comparativo de la reproducción de tres especies del género *Mobula* (Chondrichthyes: Mobulidae) en el suroeste del Golfo de California, México, Centro Interdisciplinario de Ciencias Marinas.
- Shaari Hb, Yamamoto M, Irino T (2013) Enhanced upwelling in the eastern equatorial Pacific at the last five glacial terminations. *Palaeogeography, Palaeoclimatology, Palaeoecology* **386**, 8-15.
- Shamblin BM, Bjørndal KA, Bolten AB, Hillis-Starr ZM, Lundgren I, Naro-Maciel E, Nairn CJ (2012) Mitogenomic sequences better resolve stock structure of southern Greater Caribbean green turtle rookeries. *Molecular Ecology* **21**, 2330-2340.
- Shirai S (1996) Phylogenetic interrelationships of neoselachians (Chondrichthyes: Euselachii). In: *Interrelationships of fishes* (eds. Stiassny MLJ, Parenti LR, Johnson GD). Academic Press, San Diego, CA.
- Simmons MP, Ochoterena H (2000) Gaps as characters in sequence-based phylogenetic analyses. *Systematic Biology* **49**, 369-381.
- Simpfendorfer CA, Heupel M, White WT, Dulvy NK (2011) The importance of research and public opinion to conservation management of sharks and rays: a synthesis. *Marine and Freshwater Research* **62**, 518-527.
- Slatkin M, Hudson RR (1991) Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. *Genetics* **129**, 555-562.
- Smith KA, Suthers IM (1999) Displacement of diverse ichthyoplankton assemblages by a coastal upwelling event on the Sydney shelf. *Marine Ecology Progress Series* **176**.
- Sobral AF, Afonso P (2014) Occurrence of mobulids in the Azores, central North Atlantic. *Journal of the Marine Biological Association of the United Kingdom* **94**, 1671-1675.
- Sodré D, L.F. R-F, F. SR, Rêgo PS, Schneider H, Sampaio I, Vallinoto M (2012) Inclusion of South American samples reveals new population structuring of the blacktip shark (*Carcharhinus limbatus*) in the western Atlantic. *Genetics and Molecular Biology* **35**, 752-760.
- Stamatakis A (2006) RAxML-VI-HP: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**, 2688-2690.

- Stelbrink B, von Rintelen T, Cliff G, Kriwet J (2010) Molecular systematics and global phylogeography of angel sharks (genus *Squatina*). *Molecular Phylogenetics and Evolution* **54**, 395-404.
- Stevens G (2011) Field identification guide of the prebranchial appendages (gill plates) of mobulid rays for law enforcement and trade monitoring applications.
- Stevens J (2000) The effects of fishing on sharks, rays, and chimaeras (chondrichthyans), and the implications for marine ecosystems. *ICES Journal of Marine Science* **57**, 476-494.
- Stevens JD, Bonfil R, Dulvy NK, Walker PA (2000) The effects of fishing on sharks, rays, and chimaeras (chondrichthyans), and the implications for marine ecosystems. *ICES Journal of Marine Science* **57**, 476-494.
- Stommel H, Wooster WS (1965) Reconnaissance of the Somali Current during the southwest monsoon. *Proceedings of the National Academy of Science U.S.A* **54**, 8-13.
- Strasburg JL, Rieseberg LH (2010) How robust are "Isolation with Migration" analyses to violations of the im model? A simulation study. *Molecular Biology and Evolution* **27**, 297-310.
- Sullivan J, Swofford DL, Naylor GJ (1999) The effect of taxon sampling on estimating rate heterogeneity parameters of Maximum-Likelihood models. *Molecular Biology and Evolution* **16**, 1347-1356.
- Summerer D (2009) Enabling technologies of genomic-scale sequence enrichment for targeted high-throughput sequencing. *Genomics* **94**, 363-368.
- Sumpton WD, Taylor SM, Gribble NA, McPherson G, Ham T (2011) Gear selectivity of large-mesh nets and drumlines used to catch sharks in the Queensland Shark Control Program. *African Journal of Marine Science* **33**, 37-43.
- Sun JX, Helgason A, Masson G, Ebenesersdottir SS, Li H, Mallick S, Gnerre S, Patterson N, Kong A, Reich D, Stefansson K (2012) A direct characterization of human mutation based on microsatellites. *Nature Genetics* **44**, 1161-1165.
- Suto I, Kawamura K, Hagimoto S, Teraishi A, Tanaka Y (2012) Changes in upwelling mechanisms drove the evolution of marine organisms. *Palaeogeography, Palaeoclimatology, Palaeoecology* **339-341**, 39-51.
- Swallow JC, Bruce JG (1966) Current measurements off the Somali coast during the southwest monsoon of 1964. *Deep-Sea Research* **13**, 861-888.

T

- Taguchi M, King JR, Wetklo M, Withler RE, Yokawa K (2014) Population genetic structure and demographic history of Pacific blue sharks (*Prionace glauca*) inferred from mitochondrial DNA analysis. *Marine and Freshwater Research* **66**, 267-275.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using Maximum Likelihood, evolutionary distance, and Maximum Parsimony methods. *Molecular Biology and Evolution* **28**, 2731-2739.
- Team R-C (2005) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Theisen TC, Bowen BW, Lanier W, Baldwin JD (2008) High connectivity on a global scale in the pelagic wahoo, *Acanthocybium solandri* (tuna family Scombridae). *Molecular Ecology* **17**, 4233-4247.
- Thorrold SR, Afonso P, Fontes J, Braun CD, Santos RS, Skomal GB, Berumen ML (2014) Extreme diving behaviour in devil rays links surface waters and the deep ocean. *Nature Communications* **5**, 1-7.
- Tillett BJ, Meekan MG, Broderick D, Field IC, Cliff G, Ovenden JR (2011) Pleistocene isolation, secondary introgression and restricted contemporary gene flow in the pig-eye shark, *Carcharhinus amboinensis* across northern Australia. *Conservation Genetics* **13**, 99-115.

- Tillett J, Meekan MG, Field IC, Thorburn DC, Ovenden JR (2012) Evidence for reproductive philopatry in the bull shark *Carcharhinus leucas*. *Journal of Fish Biology* **80**, 2140-2158.
- Townsend KA, Kyne PM (2010) New records of the Japanese devilray *Mobula japanica* (Muller & Henle 1841) for Australian waters. *Memoirs of the Queensland Museum* **55**, 225-230.

V

- Vaillant L, Diguët L (1898) Sur le *Cephaloptere* du Golfe de Californie. *Bulletin du Museum National d'Histoire Naturelle* **4**, 127-128.
- Vaillant LL (1879) Note sur une nouvelle espèce d'élasmodontes hypotrème, le *Cephaloptera rochebrunei*. *Bulletin de la Société Philomatique, Paris* **7**, 187-188.
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) Micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**, 535-538.
- Vignaud T, Clua E, Mourier J, Maynard J, Planes S (2013) Microsatellite analyses of blacktip reef sharks (*Carcharhinus melanopterus*) in a fragmented environment show structured clusters. *PLoS One* **8**, e61067.
- Vignaud TM, Maynard JA, Leblois R, Meekan MG, Vazquez-Juarez R, Ramirez-Macias D, Pierce SJ, Rowat D, Berumen ML, Beeravolu C, Baksay S, Planes S (2014) Genetic structure of populations of whale sharks among ocean basins and evidence for their historic rise and recent decline. *Molecular Ecology* **23**, 2590-2601.
- Vilstrup JT, Ho SY, Foote AD, Morin PA, Krieb D, Krutzen M, Parra GJ, Robertson KM, de Stephanis R, Verborgh P, Willerslev E, Orlando L, Gilbert MT (2011) Mitogenomic phylogenetic analyses of the Delphinidae with an emphasis on the Globicephalinae. *BMC Evolutionary Biology* **10**, 11-65.
- Vinas J, Bremer JRA, Pla C (2004) Inter-oceanic genetic differentiation among albacore (*Thunnus alalunga*) populations. *Marine Biology* **145**, 225-232.
- Vinayachandran PN, Yamagata T (1998) Monsoon response of the sea around Sri Lanka: generation of thermal domes and anticyclonic vortices. *Journal of Physical Oceanography* **28**, 1946-1960.

W

- Walbaum JJ (1792) Petri artemidi sueci genera piscium Röse, Grypeswaldiae, Germany.
- Walter RP, Kessel ST, Alhasan N, Fisk AT, Heath DD, Chekchak T, Klaus R, Younis M, Hill G, Jones B, Braun CD, Berumen ML, DiBattista JD, Priest MA, Hussey NE (2014) First record of living *Manta alfredi* x *Manta birostris* hybrid. *Marine Biodiversity* **44**, 1-2.
- Wandeler P, Hoeck PE, Keller LF (2007) Back to the future: museum specimens in population genetics. *Trends in Ecology and Evolution* **22**, 634-642.
- Waples RS (1987) A multispecies approach to the analysis of gene flow in marine shore fishes. *Evolution* **41**, 385-400.
- Waples RS (1998) Separating the wheat from the chaff: patterns of genetic differentiation in high gene flow species. *Journal of Heredity* **89**, 438-450.
- Waples RS, Gaggiotti O (2006) What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology* **15**, 1419-1439.
- Ward RD, Woodwark M, Skibinski DOF (1994) A comparison of genetic diversity levels in marine, freshwater and anadromous fishes. *Journal of Fish Biology* **44**, 213-232.
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PD (2005) DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **360**, 1847-1857.

- Ward-Paige CA, Davis B, Worm B (2013) Global population trends and human use patterns of *Manta* and *Mobula* rays. *PLoS One* **8**, e74835.
- Warnock J, Scherer R, Loubere P (2007) A quantitative assessment of diatom dissolution and Late Quaternary primary productivity in the Eastern Equatorial Pacific. *Deep-Sea Research Part II-Topical Studies in Oceanography* **54**, 772-783.
- Warren B, Stommel H, Swallow JC (1966) Water masses and patterns of flow in the Somali Basin during the southwest monsoon of 1964. *Deep-Sea Research* **13**, 825-860.
- Weber JL, Wong C (1993) Mutation of human short tandem repeats. *Human Molecular Genetics* **2**, 1123-1128.
- Weissing FJ, Edelaar P, van Doorn GS (2011) Adaptive speciation theory: a conceptual review. *Behavioral Ecology and Sociobiology* **65**, 461-480.
- Whitcraft S, Hofford A, Hilton P, O'Malley MP, Jaiteh V, Knights P (2014a) Evidence of declines in shark fin demand, China.
- Whitcraft S, O'Malley MP, Hilton P (2014b) The continuing threat to *Manta* and *Mobula* rays: 2013-14 market surveys, Guangzhou, China.
- White WT, Giles J, Dharmadi, Potter IC (2006) Data on the bycatch fishery and reproductive biology of mobulid rays (Myliobatiformes) in Indonesia. *Fisheries Research* **82**, 65-73.
- Whitney NM, Robbins WD, Schultz JK, Bowen BW, Holland KN (2012) Oceanic dispersal in a sedentary reef shark (*Triaenodon obesus*): genetic evidence for extensive connectivity without a pelagic larval stage. *Journal of Biogeography* **39**, 1144-1156.
- Wielstra B, Arntzen JW (2011) Unraveling the rapid radiation of crested newts (*Triturus cristatus* superspecies) using complete mitogenomic sequences. *BMC Evolutionary Biology* **11**.
- Wiley EO, Lieberman BS (2011) *Phylogenetics: theory and practice of phylogenetic systematics*, Second Edition John Wiley & Sons, Inc, Hoboken, NJ.
- Woerner AE, Cox MP, Hammer MF (2007) Recombination-filtered genomic datasets by information maximization. *Bioinformatics* **23**, 1851-1853.
- Woodward AS (1889) Catalogue of the fossil fishes in the British Museum. In: Part 1: containing the elasmobranchii (ed. Trustees of the British Museum NH).
- Woolfit M (2009) Effective population size and the rate and pattern of nucleotide substitutions. *Biology Letters* **5**, 417-420.
- Wooster WS, Reid JL (1963) *Eastern boundary currents* Wiley-Interscience, New York.
- Wooster WS, Schaefer MB, Robinson MK (1967) *Atlas of the Arabian Sea for fishery oceanography* University of California, Institute of Marine Resources, La Jolla, USA.
- Wu CI, Li WH (1985) Evidence for higher rates of nucleotide substitution in rodents than in man. *Proceedings of the National Academy of Sciences U.S.A* **82**, 1741-1745.
- Wyman SK, Jansen RK, Boore JL (2004) Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* **20**, 3252-3255.
- Wyrtki K (1961) *Physical oceanography of the southeast Asian waters* S.I.O, La Jolla, U.S.A.
- Wyrtki K (1962) The upwelling in the region between Java and Australia during the southeast monsoon. *Australian Journal of Marine Freshwater Research* **13**, 217-225.
- Wyrtki K (1964) Upwelling in the Costa Rica Dome. *Fishery Bulletin, Fish and Wildlife Service U.S.A.* **63**, 355-372.
- Wyrtki K (1966) *Oceanography of the eastern equatorial Pacific Ocean*. *Oceanography and Marine Biology, an Annual Review* **4**, 33-68.

Y

- Yu L, Li YW, Ryder OA, Zhang YP (2007) Analysis of complete mitochondrial genome sequences increases phylogenetic resolution of bears (Ursidae), a mammalian family that experienced

rapid speciation. *BMC Evolutionary Biology* **7**, 198.

Z

Zachos J, Pagani M, Sloan L, Thomas E, Billups K (2001) Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* **292**, 686-693.

Zachos JC, Quinn TM, Sallam KA (1996) High-Resolution (10(4) Years) deep-sea foraminiferal stable isotope records of the Eocene-Oligocene Climate Transition. *Paleoceanography* **11**, 251-266.

Zaytsev O, Cervantes-Duarte R, Montante O, Gallegos-Garcia A (2003) Coastal upwelling activity on the Pacific shelf of the Baja California peninsula. *Journal of Oceanography* **59**, 489-502.



Acknowledgements

As a true migratory species, I have moved between many places and across vast stretches of the open ocean while this thesis took shape. Consequently, there are many people in many corners of the world I would like to express my gratitude to. You all have made the last seven years of my life an unforgettable experience!

First of all, a huge thanks to my fearless team of PhD advisors:

Jeanine, I'm sincerely grateful for your support, especially during these last few years when it was most needed. Thank you for your guidance, tireless editing, mental support, and for helping me shape this thesis into something to be proud of.

Diego, thank you for introducing me into the magical world of mobulid research. This PhD has been a wonderful opportunity, and I am sincerely grateful for your support throughout the years, as well as your efforts to turn a geneticist into a conservationist. Also thank you and Diana for including me in your lovely family for a little while. It has been a pleasure!

Thank you Giacomo, for teaching me so many things about genetics and the world in general. I can always count on you to come up with something outrageous to change my perspective on things. Thank you also for inviting me along to Guadalupe Island for the most unforgettable spearfishing expedition ever, and the various Little Miss Sunshine-esque trips to Baja.

And last but not least: thank you Galice, for giving me the opportunity to use University of Nordland's Next Generation Sequencing (NGS) facility, and for useful discussions about data and analyses. This thesis wouldn't have been the same without you.

As with any genetic study, obtaining sufficient tissue samples is all-important. The completion of this thesis would therefore not have been possible without the help of many researchers across the world. A warm thanks to Gregor Cailliet, who -early on - provided me with a long list of his contacts to bug; Hoyt Peckham and volunteers of Proyecto Caguama, with whom I spent some hot, muggy, smelly, mosquito-infested and wonderful months in Puerto Adolfo Lopez Mateos: thank you for your help, company and (unfortunately forgotten) Spanish lessons; Thanks also to various members of the Lucero family, but especially Don Chuy from Grupo Tortugero for your help and friendship; Daniel Fernando for your help and tireless work to save mobulids - I still feel I hit the jackpot with all those samples you collected in Sri Lanka; Felipe Galvan Magaña, for providing me with samples from Baja and Ecuador; Colombo Estupiñán-Montaño for collecting samples in Ecuador; Bernard Seret, for providing me with indispensable samples from the Atlantic Ocean; Agnes Dettai for providing samples from the Muséum national d'Histoire naturelle, ichthyology collection; Robert Olson and Ernesto Altamirano from the Inter American Tropical Tuna Commission, and their observers participating in the AIDPC program, for providing samples from the East Pacific Ocean; Giuseppe Notarbartolo di Sciara and Fabrizio Serena for providing a sample of *Mobula mobular*; and finally, thank you Felix, Sanne, Annamarie and Erik for a wonderful and unforgettable sampling/disaster trip to Indonesia.

Santa Cruz has been a wonderful and unforgettable home for three* years. First of all, thank you Kelly for your warmth, friendship and all your tireless help in just about every category. If ever I discover an unclaimed rock in the middle of some tropical ocean I'll name it 'Rock Kelly'. Thanks also to all you lovely people from the Coastal Conservation Action Lab, the Bernardi Lab and the larger Long Marine Lab community: Bill, Ally, and the unforgettable Savy and Esmee, Suzy, Dena, Asha, Abe, Joe, Luz, Diana, Matthew, Erin, Angela, Yvette (2nd best spearfisherwoman in the world), Ricardo, Jimmy, Sally, Kim, Alexis, Marina, Martha, Amanda, Ana, Jose, Ana Maria, Matthieu, Dan, Ammon, Jen and Mark, Yuri, the Steinbergs, Jared and Emily, Adam, Christy and Joiah, Paul and Nancy. Finally, special thanks to Jamie, for your company, warmth, tireless support, and of course the driving lessons. California would have been a less sunny place without you!

My stay in Bodø taught me the true meaning of the phrase 'winter is coming', and I would not

have been able to survive the cold without the support and friendship of many people. Thank you Spyros, for being an amazing friend, confidant and colleague, and for the most unforgettable mushrooming expedition ever. I wish you, Marol and baby Iason all the best! Also thank you Jorge, for your friendship, lovely late-night dinners and general support, as well as for providing me with a wonderful opportunity to expand my NGS skills; Kit, for sharing your amazing warmth, home and pets with me. I'm missing our nights in front of the fire place (lots of smoke, no fire); Giulia, for providing some much needed sparkle during dark times; Andrea, for your friendship and inspiring trooper attitude; Cesilie, for those completely wasted crochet lessons; and thank you Galice, Hin, Marol, Silje, Alexander, Irina, Vigdis, Martina, Carlo, Ingvild, Boris, Christopher, Tor Erik, Åse, Chris, Teshome, Sünnje, Anne Merete, Oddvar, Tormund, Jarle, Truls and the rest of the FBA community for giving me a taste of Norway.

I have not spent much time in Groningen during my PhD, but still am indebted to a few people there. Thank you Stella for your much needed and competent help in the lab; Hinke for saving me from certain demise countless times; Corine for your warm thoughts and support; Wytze, for indispensable help with getting the thesis ready for printing; and finally my fellow Msc students, Nina, Alejo, Ivan, Michael, Ellen and Kirsten, for making science *even* more fun.

Although I've mentioned many of you already, I would specifically like to thank all the people who have provided me with a place to stay these last few years: Felix, Sanne, Lucile, Ivan, Spyros and Marol, Jorge, Kit, Andrea, Giulia, Alexander and Irina, Dena, Asha, Di and Diego, Kelly and Will, Jen and Mark, Paul, Angela, Panu, Sas and Chris. You have all made me feel extremely welcome, as well as kept me out of the cold (and covered in cat/dog hair in many cases)!

Thank you Steven Arrow, for introducing me to the wonders of the ocean through my work for Arrow Pearls. My time in Australia, Thailand and Dubai was magical, and without these experiences I would not be where I am now. Also thank you Adrian Troy (I've read all about your troubles in Eritrea mate – I loved the new haircut), and of course Fongy, to whom my thoughts still drift every now and then, wherever you might be now.

A very heartfelt thanks to friends and family who have supported me through some very difficult times these last few years. Spee, Sanne, kleine Felix, Annamarie, Manu, De (On)geboren Hoop, Sara en Jeroen: if I ever discover a new Family of mosses, I will name them after you guys. Thank you Lucile and Nina, for your warmth, friendship, advice, hospitality, long discussions about life, and Lucile for teaching me (a small part of) your cooking tricks. I miss you both a lot! Also thank you lovely people from Proeftuin de Luwte, you are a plot of fertile earth for me to thrive in. And finally, my family: Ingmar, Coby en Oma. You are a safe and warm haven to return to after this long journey!

And finally, I would like to express my sincere gratitude to my funding sources, for their faith and excellent collaboration: John 'O Sullivan and the Monterey Bay Aquarium for initiating this project and providing funding for lab consumables; Galice Hoarau and the Marine Ecology Unit, Universitet i Nordland for funding part of the Next Generation Sequencing; the Top-Master-Evolutionary Biology Program (2008–2012) of the University of Groningen, for providing a PhD stipend; the Marine Benthic Ecology and Evolution Department, CEES, University of Groningen for providing general support as well as funding for lab consumables; the Ecology and Evolutionary Biology Department, UC Santa Cruz for providing support for fieldwork; and Island Conservation for providing support for fieldwork.

Affiliations of co-authors

Bernardi, Giacomo, Department of Ecology & Evolutionary Biology, University of California Santa Cruz, 100 Shaffer Road, Santa Cruz, CA 95060, USA

Croll, Donald A., Department of Ecology & Evolutionary Biology, University of California Santa Cruz, 100 Shaffer Road, Santa Cruz, CA 95060, USA

Dewar, Heidi, Southwest Fisheries Science Centre, NOAA Fisheries, 8604 La Jolla Shore Drive, La Jolla, CA 92037, USA

Dulvy, Nicholas K., Earth to Ocean Research Group, Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6

Fernando, Daniel, Department of Biology and Environmental Science, Linnaeus University, SE 39182 Kalmar, Sweden & The Manta Trust, Catemwood House, Corscombe, Dorchester Dorset, DT2 0NT, United Kingdom

Francis, Malcolm, NIWA Taihoro Nukurangi, 301 Evans Bay Parade, Greta Point, Wellington 6021, New Zealand

Galván Magaña, Felipe, Centro Interdisciplinario de Ciencias Marinas. Instituto Politécnico Nacional. Av. IPN s/n. La Paz, Baja California Sur, Mexico. 23096

Hall, Martin, Inter-American Tropical Tuna Commission, 8901 La Jolla Shores Drive, La Jolla, CA 92037, USA

Heinrichs, Shawn, Blue Sphere Media LLC, 7382 Glacier View, Longmont, CO 80503, USA

Hoarau, Galice, Faculty of Biosciences and Aquaculture, Universitetet i Nordland, Universitetsaleen 11, 8049 Bodø, Norway

Kollias, Spyridon, Centre for Ecological and Evolutionary Synthesis, Universitet i Oslo, Blindernveien 31, 0316 Oslo, Norway

Marshall, Andrea, Marine Megafauna Foundation, 11260 Donner Pass Rd, 256 Truckee, CA 96161, USA

McCauley, Douglas, Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, CA 93106, USA

Newton, Kelly M., Department of Ecology & Evolutionary Biology, University of California Santa Cruz, 100 Shaffer Road, Santa Cruz, CA 95060, USA

Notarbartolo di Sciara, Giuseppe, Tethys Research Institute, Acquario Civico, 20121 Milano, Italy

Olsen, Jeanine L., Department of Marine Benthic Ecology and Evolution, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands

O'Malley, Mary, The Manta Trust, Catemwood House, Corscombe, Dorchester Dorset, DT2 0NT, United Kingdom & Wildaid, 744 Montgomery Street, Suite 300, San Francisco, CA 94111

O'Sullivan, John, Monterey Bay Aquarium, 886 Cannery Row, Monterey, CA 93940, USA

Roman, Marlon, Inter-American Tropical Tuna Commission, 8901 La Jolla Shores Drive, La Jolla, CA 92037, USA

Seret, Bernard, Muséum national d'Histoire naturelle, Département Systématique et Evolution, CP 51, rue Buffon, 75231 Paris cedex 05, France

Smolina, Irina, Faculty of Biosciences and Aquaculture, Universitetet i Nordland, Universitetsaleen 11, 8049 Bodø, Norway

Stevens, Guy, The Manta Trust, Catemwood House, Corscombe, Dorchester Dorset, DT2 0NT, United Kingdom

Tershy, Bernie R., Department of Ecology & Evolutionary Biology, University of California Santa Cruz, 100 Shaffer Road, Santa Cruz, CA 95060, USA

Wintner, Sabine, KwaZulu-Natal Sharks Board, 1A Herrwood Drive, Umhlanga Rocks 4320, South Africa & Biomedical Resource Unit, University of KwaZulu-Natal, Westville Campus, University Road, Westville 3600, South Africa

List of publications

Published

Kollias S, **Poortvliet M**, Smolina I, Hoarau G (2015) Low cost sequencing of mitogenomes from museum samples using baits capture and Ion Torrent. *Conservation Genetics Resources* **7**, 345-348.

Poortvliet M, Olsen JL, Croll DA, Bernardi G, Newton KM, Kollias S, O'Sullivan J, Fernando D, Stevens G, Galván Magaña F, Seret B, Wintner S, Hoarau G (2015) A dated molecular phylogeny of manta and devil rays (Mobulidae) based on mitogenome and nuclear sequences. *Molecular Phylogenetics and Evolution* **83**, 72- 85.

Smolina, I, Kollias S, **Poortvliet M**, Nielsen TG, Lindequ P, Castellani C, Møller EF, Blanco-Bercial L, Hoarau G (2014) Genome- and transcriptome-assisted development of nuclear insertion/deletion markers for *Calanus* species (Copepoda: Calanoida) identification. *Molecular Ecology Resources* **14**, 1072- 1079.

Poortvliet M, Longo GC, Selkoe K, Barber PH, White C, Caselle JC, Perez Matus A, Gaines SD, Bernardi G (2013). Phylogeography of the California sheephead, *Semicossyphus pulcher*: the role of deep reefs as stepping-stones and pathways to antitropicality. *Ecology and Evolution* **3**, 4558-71.

Poortvliet M, Hoarau G (2013) The complete mitochondrial genome of the Spinetail Devilray, *Mobula japanica*. *Mitochondrial DNA* **1**, 28-30.

Poortvliet M, Galván Magaña F, Bernardi G, Croll DA, Olsen JL (2011) Isolation and characterization of twelve microsatellite loci for the Japanese Devilray (*Mobula japanica*). *Conservation Genetics Resources* **3**, 733-35.

Poortvliet M, Olsen JL, Selkoe KA, Coyer JA, Bernardi G (2009) Isolation and characterization of 11 microsatellite primers for a temperate reef fish, the California sheephead (*Semicossyphus pulcher*). *Molecular Ecology Resources* **9**, 429-30.

Submitted

Croll DA, Dewar H, Dulvy NK, Fernando D, Francis MP, Galván-Magaña F, Hall M, Heinrichs S, Marshall A, McCauley D, Newton KM, Notarbartolo-di-Sciara G, O'Malley M, O'Sullivan J, **Poortvliet M**, Roman M, Stevens G, Tershy BR, White WT Vulnerabilities and Fisheries Impacts: The Uncertain Future of Manta and Devil Rays. Submitted to Aquatic Conservation

In preparation

Poortvliet M, Hoarau G, Croll DA, Bernardi G, Newton KM, O'Sullivan J, Fernando D, Stevens G, Galván Magaña F, Seret B, Olsen JL Influences of past climatic change on historical population structure and demography of the Spinetail devil ray, *Mobula japanica*, based on mitogenomes and nuclear DNA. In preparation for *Molecular Ecology*

